



FIRE-RES

Innovative technologies & socio-ecological-economic solutions for fire resilient territories in Europe

D1.12 Innovative post-fire strategies and adaptation to the current context of increasing environmental uncertainties

www.fire-res.eu

fire-res@ctfc.cat

Project Acronym: FIRE-RES

Project name: Innovative technologies and socio-ecological-economic solutions for fire resilient territories in Europe

Call ID: H2020-LC-GD-1-1-2020 (Preventing and fighting extreme wildfires with the integration and demonstration of innovative means)

Work Package: 1

Task Number: 1.4.2

Lead beneficiary: Forest Science and Technology Centre of Catalonia (CTFC)

Contributing beneficiary(ies): Catalan Fire and Rescue Service (CFRS), Instituto Superior de Agronomía (ISA), Forest Owners' Association of the Sousa Valley (AVFS), University of Forestry of Sofia (US), Corporación Nacional Forestal (CONAF), European Forest Institute (EFI)



This document was produced under the terms and conditions of Grant Agreement No. 101037419 of the European Commission. It does not necessarily reflect the view of the European Union and in no way anticipates the Commission's future policy in this area.

Publication

Publication date: 31/05/2023

Authors: Teresa Valor, Pere Casals, Míriam Piqué, Lluís Coll

Abstract: This document is the first of two deliverables that focus on post-fire strategies and adaptation to increasing environmental uncertainties. Its goal is to identify high-priority areas for post-fire restoration across different forest types. The report draws on a literature review and survey of wildfire experts to identify key factors, metrics, and thresholds that determine post-fire recovery rates. Factors such as pre-fire vegetation, fire regime, topography, soil, and pre/post-fire climate were identified as the main drivers of post-fire vegetation recovery. The report provides a tabular summary of factors and associated thresholds to predict vulnerable areas at higher risk of erosion and runoff both before and in the short-term after the fire across different forest types.

Keywords: post-disturbance dynamics, resilience, fire adaptation, restoration, fire severity, plant strategies

Quote as: Valor, T; Casals, P; Pique, M; Coll, L; (2023). Innovative post-fire strategies and adaptation to the current context of increasing environmental uncertainties Deliverable D1.12 FIRE-RES project. 56 pages. DOI: 10.5281/zenodo.7966120

DOI: 10.5281/zenodo.7966120

Dissemination level

[x] PU- Public: must be available in the website

[] CO- Confidential: Only for members of the Consortium and the Commission Services

[] CI – Classified: As referred in to Commission Decision 2001/844/EC

Document history

Edition	Date	Status	Author
Version 1	15/04/2023	Draft	Teresa Valor (CTFC), Pere Casals (CTFC), Míriam Piqué (CTFC), Lluís Coll (CTFC)
Version 2	16/05/2023	Revision	Edgar Nebot (CFRS)
Version 3	21/05/2023	Final version	Teresa Valor (CTFC), Pere Casals (CTFC), Míriam Piqué (CTFC), Lluís Coll (CTFC)

Copyright © All rights reserved. This document or any part thereof may not be made public or disclosed, copied or otherwise reproduced or used in any form or by any means, without prior permission in writing from the FIRE-RES Consortium. Neither the FIRE-RES Consortium nor any of its members, their officers, employees or agents shall be liable or responsible, in negligence or otherwise, for any loss, damage or expense whatever sustained by any person as a result of the use, in any manner or form, of any knowledge, information or data contained in this document, or due to any inaccuracy, omission or error therein contained.

All Intellectual Property Rights, know-how and information provided by and/or arising from this document, such as designs, documentation, as well as preparatory material in that regard, is and shall remain the exclusive property of the FIRE-RES Consortium and any of its members or its licensors. Nothing contained in this document shall give, or shall be construed as giving, any right, title, ownership, interest, license or any other right in or to any IP, know-how and information.

The information and views set out in this publication does not necessarily reflect the official opinion of the European Commission. Neither the European Union institutions and bodies nor any person acting on their behalf, may be held responsible for the use which may be made of the information contained therein.

Table of Contents

1	INTRODUCTION	1
2	DELIVERABLE AIMS	1
3	METHODOLOGY	2
3.1	LITERATURE REVIEW	2
3.2	QUESTIONNAIRE ON FACTORS AFFECTING RESISTANT AND RESILIENT STAND AND LANDSCAPES.....	3
4	REVIEW ON FACTORS MODULATING FIRE IMPACTS AND VEGETATION RECOVER	3
4.1	FIRE INTOLERANT SPECIES AND FIRE REGIME FACTORS	7
4.1.1	<i>Pre fire vegetation: species traits and individual, stand and landscape characteristics.</i>	7
4.1.2	<i>Fire severity</i>	8
4.1.3	<i>Fire frequency</i>	8
4.1.4	<i>Fire season</i>	8
4.2	FIRE RESISTANT SPECIES AND FIRE REGIME FACTORS.....	8
4.2.1	<i>Pre fire vegetation: species traits and individual, stand and landscape characteristics.</i>	8
4.2.2	<i>Fire severity</i>	9
4.2.3	<i>Fire frequency</i>	11
4.2.4	<i>Fire season</i>	11
4.3	FIRE SENSITIVE SPECIES AND FIRE REGIME FACTORS.....	12
4.3.1	<i>Pre-fire vegetation: species traits and individual, stand and landscape characteristics.</i>	12
4.3.2	<i>Fire severity</i>	13
4.3.3	<i>Fire frequency</i>	14
4.3.4	<i>Fire season</i>	14
4.4	FIRE TOLERANT SPECIES AND FIRE REGIME FACTORS	15
4.4.1	<i>Pre fire vegetation: species traits and individual, stand and landscape characteristics.</i>	15
4.4.2	<i>Fire severity</i>	16
4.4.3	<i>Fire frequency</i>	16
4.4.4	<i>Fire season</i>	16
4.5	TOPOGRAPHY	17
4.6	SOIL CHARACTERISTICS.....	18
4.7	PRE-FIRE AND POST-FIRE CLIMATIC CONDITIONS	19
4.8	OTHER FACTORS.....	20
5	QUESTIONNAIRE RESULTS ON FACTORS DRIVING POST-FIRE DYNAMICS	20
6	SUMMARY: CRITERIA TO IDENTIFY VULNERABLE AREAS FOR RESTORATION BEFORE AND AFTER FIRE OCCURRENCE	28
7	REFERENCES	31
8	APPENDICES	40
8.1	PRELIMINARY DOMINANT SPECIES SELECTED.....	41
8.2	LITERATURE REVIEW: SEARCH STRATEGY	45
8.3	QUESTIONNAIRE	46

List of figures

Figure 1. Framework to plan post-fire management and restoration in burned area (modify from Moreira et al., 2012). The focus of the current document is predicting vulnerable areas based on pre-fire vegetation type, fire regime, soil characteristics, topography, climatic conditions and landscape structure before and short-term after fire occurrence.....	2
Figure 2. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of hemi boreal and continental Scots pine (<i>P. sylvestris</i>) forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.....	21
Figure 3. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of alpine <i>P. sylvestris</i> or <i>P. nigra</i> in the Alps or <i>P. uncinata</i> in the Pyrenees. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.....	22
Figure 4. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of spruce (<i>Picea abies</i>). The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.	23
Figure 5. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of tall deciduous oak (<i>Quercus</i> sp.) forest. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.	24
Figure 6. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of mixed <i>Quercus</i> sp. and <i>Fraxinus</i> forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.	25
Figure 7. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of fir <i>Abies alba</i> forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.	26
Figure 8. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of beech (<i>Fagus</i> sp.) forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.	27

List of tables

Table 1. Post-fire strategies of selected dominant overstory species in a range of European forest types (see Appendices: Table 1). The assignment of post-fire regeneration strategy to each species was done using the database BROT 2.0. If the database BROT 2.0 did not contain data on fire-related traits for a given species, a literature search was carried out to determine the post-fire regeneration strategy. In such cases, the reference is given as a note in the table.	5
Table 2. Fire resistant, sensitive, and tolerant European forest types and associated dominant species.	6
Table 4. Preliminary selection of species for conducting the literature review on the influence of different factors on post-fire recovery. The selection of species was based on the work of Pividori et al., (2016), who classified the species presence for each European forest type (EFT) of the EFT classification into three classes: the species is abundant and dominant in the EFT; the species presence in the EFT is either secondary or predominant, but under particular and uncharacteristic ecological conditions of the EFT; the presence in the EFT is in some cases both dominant and secondary. Only species that were abundant and dominant in a given EFT were selected from these categories.	41
Table 5. Keyword searches for each fuel-related factors associated with stand and landscape resistance.	45

1 Introduction

The overall goal of this report is to enhance our comprehension of the variables influencing the ability of forests to respond to (and recover from) fires. This information holds significant importance in assessing the ecological resilience of the landscape. Ecological resilience is the ability of the ecological system to regain its previous functions and ecosystem services following a fire incident. Previous studies have shown that post-fire vegetation recovery is influenced by a number of factors related to pre-fire vegetation (e.g., forest composition and structure), fire regime (fire severity, fire frequency, fire season), soil properties (stoniness, lithology...), topography (slope, orientation), landscape structure (forest cover, diversity and distribution of land use) and climatic conditions before and after the fire event.

This report relies on the results derived from research conducted on wildfire and prescribed burning experiments of different intensities and severities. Emphasis was placed on studies conducted in new fire areas, since EWEs do not necessarily lead to ecological disasters (the inherent unpredictable behaviour of EWE can lead to a mosaic of different severities).

2 Deliverable aims

The primary objective of this document is to compile and integrate existing information concerning the main factors driving the recovery of vegetation in fire-prone and new fire areas. By utilizing this information, our aim is to facilitate the establishment of science-based criteria for identifying priority post-fire recovery areas, i.e. defining ecological units (e.g. species, communities...) that may be vulnerable to fire and might require management intervention if burned. The type of interventions required will be discussed later in *D. 12: Report on innovative post-fire strategies and adaptation to the current context of increasing environmental uncertainties*, scheduled for month 40 of the FIRE-RES project. Furthermore, this report provides the essential information for developing implementation guidelines for the *IA. 2.6 brief: Designing post-fire restoration strategies*. Consequently, the principal outcome is a summary table that outlines key factors, metrics, and thresholds for predicting vulnerable areas before and just after a fire occurrence.

This document presents essential information for conducting the initial step of the framework developed by Moreira et al., (2012) to plan post-fire management and restoration in burned areas. The process entails the identification of priority areas for post-fire interventions both before and after a fire event (Figure 1). To accurately predict areas that are vulnerable to fire before it occurs, it is imperative to establish thresholds for various factors such as vegetation, soil characteristics, topography, landscape structure, and pre-fire climatic conditions. Conversely, to identify vulnerable areas shortly after a fire, pre-fire thresholds need to be combined with thresholds associated with factors such as the characteristics of the fire itself (i.e., the fire regime), post-fire conditions, competition with pioneer

species, and pest outbreaks. This comprehensive approach facilitates a more effective and targeted intervention strategy.



Figure 1. Framework to plan post-fire management and restoration in burned area (modified from Moreira et al., 2012). The focus of the current document is predicting vulnerable areas based on pre-fire vegetation type, fire regime, soil characteristics, topography, climatic conditions and landscape structure before and short-term after fire occurrence.

3 Methodology

The information presented in this deliverable was gathered by employing two methods: 1) a systematic literature review of the factors, metrics and associated thresholds affecting post-fire recovery for a range of forest types, and 2) a questionnaire distributed to FIRE-RES participants, foresters and forest scientists. The main output of this research is a tabular summary with a list of species (that are linked to the European Forest Types) classified in terms of their post-fire regeneration strategies and the identified factors and associated thresholds affecting their recovery. This summary may allow the identification of priority areas for establishing post-fire restoration actions for each forest type. More detailed information regarding the specific methodologies employed can be found in Appendix 8.1 (Literature review: search strategy) and 8.2 (Questionnaire).

3.1 Literature review

A preliminary selection of forest types from the European Forest Type Classification System (EFT) was carried out based on the work of Pividori et al.,

(2016), who classified overstory species for each European forest type (EFT) depending on their abundance and dominance. Only those species that exhibited significant abundance and dominance in a given EFT were selected to conduct the literature review on the factors influencing their recovery after fire (Appendices: Table 1). In addition to the overstory species, heathlands dominated by *Calluna vulgaris* were included because they are important habitats in northern Europe, and *Quercus coccifera* garrigues due to their importance in Mediterranean countries.

3.2 Questionnaire on factors affecting resistant and resilient stand and landscapes

The results of the literature review were used to formulate a questionnaire concerning the factors influencing post-fire dynamics for specific forest types. The aim of the questionnaire was as follows:

- To collect specialized knowledge regarding the various factors that impact the dynamics of some vegetation types from boreal, alpine and continental bioregions after fires, where information is currently lacking. This is important to make recommendations and set priorities for post-fire management in those particular forest types.

The questionnaire was distributed to experts in the field of forest fires in Europe and Chile. Section 5 presents the results of the questionnaire. The questionnaire also included questions on the factors and thresholds that determine the resilience and resistance of landscapes to EWE. These factors were discussed in *D 1.11 Ecological factors driving resistant and resilient landscapes to high intensity and extreme wildfire events* (Valor et al., 2023).

4 Review on factors modulating fire impacts and vegetation recover

Plants can be classified depending on their post-fire regeneration strategies as obligate resprouters, facultative seeders/resprouters, obligate seeders, and fire colonizers depending on their post-fire regeneration strategy (*Box 1*, see Pausas & Keeley, 2014).

Box 1. Post-fire regeneration strategies as defined in Pausas & Keeley (2014). As noted by the authors, the terms "seeders" and "resprouters" refer exclusively to post-fire conditions and cannot be used to plants that regenerate by seed or resprout under other conditions.

Obligate resprouters (R+S-): plants that rely on resprouting to regenerate after fire (resprouters without postfire seeding ability). These plants do not germinate after fire because they lack a fire-resistant seed bank.

Obligate seeders (R-S+): plants that do not resprout and rely on seeding to regenerate their population after fire (nonresprouters with postfire seeding ability).

Facultative seeders/resprouters (R+S+): plants that have both mechanisms for regenerating after fire, that is, they are able to resprout and to germinate after fire.

Post-fire colonizers (R-S-): plants that lack a mechanism for local postfire persistence, but they recruit after fire by seeds dispersed from unburned patches or from populations outside the fire perimeter (metapopulation dynamics).

Priority areas for post-fire restoration are generally those showing slow plant recovery and high risk of erosion and runoff after fire (Vallejo et al., 2012). The rate of vegetation recovery after a fire varies among forest types and depends on the presence and abundance of fire-adapted plant regeneration strategies in the system (Vallejo et al., 2012). Generally, a variety of fire-adapted plant characteristics coexist within a forest type, but the recovery of vegetation communities is often assessed based on the response of the most readily identifiable vegetation or the species dominating the overstory (Nolan et al., 2021). In this study, our focus is directed towards analyzing the factors driving the capacity of overstory species to recover or survive after a fire. To accomplish this, we utilized the BROT 2.0 database developed by Tavşanoğlu & Pausas (2018), fire-related traits (e.g., fire-stimulated flowering, ability to resprout after fire, heat-stimulated germination, seedling emergence after fire, fire-related chemical cues, thick bark, canopy seed bank) were compiled for the preliminary selected dominant species (Appendices: Table 3). However, for certain species, fire-related trait data was not available in the BROT 2.0 database. In such instances, we searched for their post-fire regeneration strategies in scientific articles using the species name and "post-fire regeneration" as keywords. The species that lacked data in both the BROT 2.0 database and the ISI Web of Knowledge were removed from the list and not considered for the literature review.

Table 1. Post-fire strategies of selected dominant overstory species in a range of European forest types (see Appendices: Table 3). The assignment of post-fire regeneration strategy to each species was done using the database BROT 2.0. If the database BROT 2.0 did not contain data on fire-related traits for a given species, a literature search was carried out to determine the post-fire regeneration strategy. In such cases, the reference is given as a note in the table.

Obligate resprouters	Obligate seeder		Post-fire colonizer	No fire-related traits
	Soil or canopy seed bank	No soil or canopy seed bank		
<i>Pinus canariensis</i>	<i>Pinus pinaster</i>	<i>Pinus pinea</i>	<i>Pinus pinea</i>	<i>Abies alba</i> ¹
<i>Quercus suber</i>	<i>Pinus halepensis</i>	<i>Pinus nigra</i>	<i>Pinus nigra</i>	<i>Picea abies</i> ¹
<i>Quercus ilex</i>	<i>Calluna vulgaris</i> ³	<i>Pinus sylvestris</i>	<i>Pinus sylvestris</i>	
<i>Quercus robur</i>		<i>Pinus mugo</i> ¹	<i>Pinus mugo</i> ¹	
<i>Quercus pubescens</i>		<i>Pinus cembra</i> ²	<i>Pinus cembra</i> ²	
<i>Fraxinus angustifolia</i>		<i>Larix decidua</i> ²	<i>Larix decidua</i> ²	
<i>Fagus sylvatica</i>				
<i>Castanea sativa</i>				
<i>Populus tremula</i>				
<i>Betula pendula</i>				
<i>Quercus coccifera</i>				
<i>Calluna vulgaris</i> ³				

¹ Feurdean et al., (2019)

² Frejaville et al., (2013)

³ *Calluna vulgaris* can either resprout or germinate after fire.

The identification of the post-fire strategy of the dominant species allowed to classify the selected forest types as fire-tolerant (dominated by resprouters species), fire sensitive (dominated by obligate seeders), fire resistant (dominated by post-fire colonizers species), or fire intolerant (dominated by species with no fire-related traits that can re-colonize burned areas from adjacent unburned patches) (Table 2).

Table 2. Fire resistant, sensitive, and tolerant European forest types and associated dominant species.

Fire type forest	Dominant species	European Forest type
Fire intolerant	<i>Abies alba</i>	2.8 Nemoral silver fir
		10.6 Mediterranean and Anatolian fir forest
	<i>Picea abies</i>	1.1 Spruce and spruce-birch boreal forest
		2.1 Hemiboreal forest
		2.3 Nemoral spruce forest
		3.2 Subalpine and mountainous spruce and mountainous mixed spruce-silver fir forest
	11.1 Spruce mire forest	
Fire resistant	<i>Pinus nigra</i>	2.4 Nemoral Black pine forest
		10.2.1 Mediterranean Black pine forest
		10.2.2 Anatolian Black pine forest
	<i>Pinus sylvestris</i>	1.2 Pine and pine-birch boreal forest
		2.1 Hemiboreal forest
		2.2 Nemoral Scots pine forest
		2.5 Mixed Scots pine-birch forest
		10.4 Mediterranean and Anatolian Scots pine forest
		11.2 Pine mire forest
	<i>Pinus pinea</i>	10.1.3 Mediterranean pine forest - <i>Pinus pinea</i>
<i>Pinus mugo</i>	3.1 Subalpine larch-arolla pine and dwarf pine forest	
<i>Pinus cembra</i>	3.1 Subalpine larch-arolla pine and dwarf pine forest	
<i>Larix decidua</i>	3.1 Subalpine larch-arolla pine and dwarf pine forest	
Fire sensitive	<i>Pinus halepensis</i>	10.1.2 Mediterranean pine forest - <i>Pinus halepensis</i>
	<i>Pinus pinaster</i>	2.7 Atlantic maritime pine forest
		10.1.1 Mediterranean pine forest - <i>Pinus pinaster</i>
	<i>Calluna vulgaris</i>	Heathlands
Fire tolerant	<i>Pinus canariensis</i>	10.3 Canarian pine forest
	<i>Populus tremula</i>	11.6 Aspen swamp forest
		13.4 Aspen forest
	<i>Fagus sylvatica</i>	6.1 Lowland beech forest of southern Scandinavia and north central Europe
		6.2 Atlantic and subatlantic lowland beech forest
		6.3 Subatlantic submountainous beech forest
		6.4 Central European submountainous beech forest
		6.5 Carpathian submountainous beech forest
		6.6 Illyrian submountainous beech forest
		6.7 Moesian submountainous beech forest
		7.1 South western European mountainous beech forest
7.2 Central European mountainous beech forest		
7.3 Apennine-Corsican mountainous beech forest		

	7.4 Illyrian mountainous beech forest
	7.5 Carpathian mountainous beech forest
	7.6 Moesian mountainous beech forest
<i>Castanea sativa</i>	8.7 Chestnut forest
<i>Quercus coccifera</i>	9.1 Mediterranean evergreen oak forest
<i>Quercus ilex</i>	9.1 Mediterranean evergreen oak forest
<i>Quercus suber</i>	9.1 Mediterranean evergreen oak forest
<i>Quercus robur</i>	11.5 Pedunculate oak swamp forest
<i>Quercus pubescens</i>	8.1.1 Downy oak forest - western
	8.1.2 Downy oak forest - Italian
	8.1.3 Downy oak forest - Greek, Anatolian
	8.1.4 Downy oak forest - steppe
<i>Fraxinus angustifolia</i>	8.8.1 Thermophilous ash forest
<i>Betula pendula</i>	3.4 Mountainous birch forest
	13.3 Birch forest
<i>Calluna vulgaris</i>	Heathlands

At each fire forest type level, we proceed to examine specific studies in the subsequent sections that have investigated the impact of pre-fire vegetation characteristics (species-specific fire-related traits) as well as variables at the individual (e.g. age, DBH) or stand/landscape level (e.g. canopy cover). Additionally, we also investigated associated thresholds that enable them to interact appropriately with their reference fire regime (severity, frequency, fire season). Furthermore, when accessible, we present data on the recuperation or resilience of the predominant species or fire-type forest in relation to topographic and soil characteristics, as well as pre- and post-fire conditions.

4.1 Fire intolerant species and fire regime factors

4.1.1 Pre fire vegetation: species traits and individual, stand and landscape characteristics.

Fire-intolerant species are not adapted to any fire regime as these species lack of fire-related traits. Such is the case with *P. abies* and *A. alba*, both distinguished by their thin bark and the absence of a canopy or soil seed bank (Rogers et al., 2015). In fact, Bär & Mayr, (2020) discovered that the bark of *P. abies* provides the least thermal insulation for the internal tissues of the ten central alpine tree species studied. In the same study, *A. alba* exhibited a higher insulating capacity than *P. abies*, albeit with limited overall cambium thermal protection. Therefore, *P. abies* and *A. alba* must be classified as fire intolerant, given their high susceptibility to crown and surface fires. However, a DBH above 20 cm may lead to lower mortality after fire (Dupire et al., 2019) and, thus to a higher probability of regeneration from slightly affected trees.

4.1.2 Fire severity

No studies were found on the effect of fire severity (high or low) on post-fire regeneration or resistance to fire of fire-intolerant species.

4.1.3 Fire frequency

There is a scarcity of research concerning the impacts of fire frequency on *P. abies* and *A. alba*. For *P. abies*, studies based on macrofossil and pollen analyses have shown that the abundance of this species remained stable under moderately severe and frequent fires during the Holocene in the Czech Republic (Carter et al., 2018) and in the Carpathians with fire frequencies of 200–300 years (Feurdean et al., 2017) and of 1000–4000 years (Finsinger et al., 2018). A recent study has shown that regeneration of *P. abies* was relatively low during the first 10 years after a surface fire in the Arkhangelsk region (Russia) but gradually increased the years after (Ananyev et al., 2022). In this particular case, fire severity did not influence post-fire regeneration abundance. Regarding *A. alba*, the paleoecological study of Tinner et al., (2000) revealed that high fire severity can lead to its local extinction. More recently, a study based on mid-Holocene data suggested that *P. abies* and *A. alba* forests may be most vulnerable at increased fire frequency and intensity (Feurdean et al., 2019). The authors also conclude that *A. alba* shows some resilience at intermediate fire frequency.

4.1.4 Fire season

Studies analysing the effect of fire season on the resistance or recovery of the selected dominant fire intolerant species are lacking. However, the effect of fire season on the potential tree mortality of both species has been analysed in a fire simulation study (Dupire et al., 2019). Results show that low mortality can be expected for both species after fires in the cold season but under drier conditions, mortality can reach values above 50%, especially in young forests (DBH < 20cm).

4.2 Fire resistant species and fire regime factors

4.2.1 Pre fire vegetation: species traits and individual, stand and landscape characteristics.

Fire regimes suitable for forests dominated by fire-resistant species are characterised by a combination of low intensity and frequent wildfires (Pausas, 2015). The majority of the selected fire-resistant species (*P. nigra*, *P. sylvestris*, *P. pinea*, *P. cembra*, *P. mugo* and *L. decidua*) possess a thick bark when mature and some have the ability to self-pruning, enabling them to safeguard sensitive tissues from fire. These traits not only allow their survival but also facilitate recolonization of the fire-affected open spaces.

The degree of fire resistance of the species varies. According to Fernandes et al., (2008) the ranking of fire resistance (from highest to lowest) for the most represented pine species would be: *P. pinea* > *P. nigra* > *P. sylvestris*. For a tree of 15 cm of DBH and 8 m height, the time required at temperatures above 60° C (at fire intensity of 400 kW m⁻¹) to kill their cambium would be 6.5, 5.4 and 3.7

minutes, respectively (Fernandes et al., 2008). For the other selected species, Bär & Mayr, (2020) ranked the resistance of *L. decidua* and *P. cembra* to cambium death after *P. sylvestris* (with *P. cembra* showing higher resistance than *L. decidua*). In *P. cembra*, bark samples thicker than 8 mm showed a marked increase in heat isolation potential. As noted by Bär & Mayr, (2020), this species has the thinnest bark within the genus. Finally, *P. mugo* is considered a fire-resistant species as its branches retain moisture from precipitation and snowmelt under the canopy. In return, it can promote fire spread due to its layering effect that increases fuel connectivity (Leys et al., 2014).

Few studies have addressed the effect of pre-fire stand or landscape variables fire on the post-fire resistance or regeneration of the selected species. In the case of *P. nigra*, Martín-Alcón & Coll (2016) found higher post-fire regeneration in sites with a long history of tree cover and high pre-fire tree cover (> 50%). Moreover, post-fire regeneration was mainly determined by the abundance of unburned patches after the fire (Martín-Alcón & Coll, 2016; Ordóñez et al., 2006). Therefore, the presence of mature pre-fire stands with low fuel loads and high vertical discontinuity is an important factor for ensuring the post-fire regeneration of resistant species. In another study, pre-fire land use history, particularly the time since cropland abandonment, was examined as a potential factor influencing the abundance of different species after the fire (Puerta-Piñero et al., 2012). The authors concluded that time since cropland abandonment did not appear to influence the abundance of *P. nigra* and *P. sylvestris*. Regarding understory composition, Vallejo et al. (2012) pointed out that the recovery of forests dominated by obligate seeders mostly depends on the composition of the understory, as pines take longer to establish. In these cases, burnt areas previously dominated by a forest with a resprouting understory may have lower erosion risk than those dominated by obligate seeders.

4.2.2 Fire severity

The interaction between fire severity, species traits and the age or size of individuals determine the ability of a fire-resistant species to survive a fire. Damage to the cambium by fire depends on the fire characteristics (e.g., temperature and residence time) as well as the tree's ability to protect sensitive tissues. The ability to protect tissues and survive depends on the species identity and the bark thickness for a given tree size (Catry et al., 2010). In low to moderate intensity fires, adult fire-resistant species usually survive because their bark is thick enough to withstand such intensities. However, at high intensity fires survival depends on the degree of fire damage suffered not only to the cambium but also to the crown. If tree mortality occurs, post-fire recovery may be limited because these species lack fire-related traits for massive post-fire recruitment.

Low severity fires

For most fire-resistant species, tree size thresholds that determine survival after low-intensity fires have been defined. In Central Catalonia (NE Spain), *P. nigra* trees with a DBH between 7.5-20 cm and a bole scorched height of 3 m were able to

survive a surface fire of moderate intensity (Valor et al., 2013). Furthermore, fire simulations show that low mortality can be expected on stands with *P. nigra* individuals with a DBH > of 20 cm (Dupire et al., 2019).

Tree size also plays a critical role in the resistance of *P. sylvestris* to low intensity fires. After a moderate-intensity experimental fire with bole scorch height of up to 2 m, the probability of tree mortality of this species was below 20% for trees with a DBH > 20 cm (Valor et al., 2017). In fact, the DBH threshold of 20 cm for low to moderate intensity fires seems to be quite consistent across different studies conducted in *P. sylvestris* (see Dupire et al., (2019) for the Alps, Linder et al., (1998) in northern Sweden and Sidoroff et al., (2007) in southern Finland). In the case of *P. pinea*, Madrigal et al., (2019) established a critical threshold for bark thickness of 2 cm. Below this value, the rate of heat transfer increases, reducing the time to reach lethal temperatures in the cambium and thus the resistance to fire. Similarly, *P. pinea* individuals with a DBH of 40 ± 11 cm and mean bark thickness at the base of the trunk of 3.1 ± 0.75 cm did not suffer mechanical damage to stems, nor significant reduction in radial growth after a prescribed burning experiment in Italy (flame length at the base of the trunk never exceeded 1.5 m) (Battipaglia et al., 2016). For *P. cembra*, *P. mugo* and *L. decidua* the lack of available data on low intensity fires does not allow us to provide tree size thresholds.

Moderate and high severity fires

Under conditions of long fire-free intervals, fire-resistant species can succumb to fire even when they are mature because fuel accumulation can lead to high fire intensity/severity. Vulnerability to crown kill is highest in *P. sylvestris* followed by *P. nigra* and then by *P. pinea* (Fernandes et al., 2008). Bär et al., (2021) showed that closed buds of *P. cembra* can withstand the highest exposure temperatures after *P. sylvestris* while *L. decidua* would be the least resistant (no data for *P. mugo* exists).

Post-fire regeneration of fire-resistant species affected by crown fires is dependent on the survival of individuals or the presence of unburnt patches from which seeds can be dispersed into the completed burned area. For instance, the high intensity wildfire that occurred in Central Catalonia in 1998 burned primarily mature *P. nigra* stands whose understory consisted mainly of oaks (Martín-Alcón & Coll, 2016). Today, these areas are mainly covered with young oaks. Regeneration of pines occurs in areas near the fire site where unburned pines have scattered their seeds into the fire area (Ordóñez et al., 2006), and within the fire perimeter around black pine islands that were not burned (Martín-Alcón & Coll, 2016). In contrast, in the case of *P. pinea*, Rodrigo et al., (2007) show that post-fire regeneration was very low in 8 areas affected by forest fires in Catalonia, due to the short dispersal distance of seeds of this species and the low survival rate of seedlings. While *L. decidua* has a relatively thick bark and is therefore considered a fire-resistant species, it has been shown to have strong recruitment after fires of moderate severity. When large areas are affected at high intensity, larch regeneration may be limited (Moris et al., 2017b).

Crown scorch thresholds that determine the probability of post-fire mortality for some of the selected species have been derived from modelling studies. For example, in the margins and in the vicinity of the islands of vegetation that survived a high-intensity fire occurred in 1998 in NE Spain, *P. nigra* individuals were alive when the extent of crown scorched was lower than 1/3. When the proportion of crown scorched varied between 1/3 to 2/3 almost 80% of the individuals survived, and more than 60% did it for crown scorched ratios above 2/3 (Ordóñez et al., 2005). It is worth mentioning that there might be differences in the effect of crown scorch on tree mortality depending on the fire season, with spring scorched being more damaging to the tree than autumn and winter fires. For example, in the case of *P. nigra* and *P. sylvestris*, trees with more than 80% of the damaged crown died one year after a spring fire, but 20% survived after an autumn fire (Valor et al., 2017). Similarly, tree survival three years after a winter fire in France was 91% (*P. nigra*) and 70% (*P. sylvestris*) for both species with less than 2/3 of the crown completely burnt (Rigolot data in Fernandes et al., 2008). In the case of *P. pinea*, about 90% of the individuals were most likely to die when the mean proportion of charred stem length was above 60%, and most likely to survive when it was below 30% (Rigolot, 2004). Similarly, Catry et al., (2010) showed that survival of *P. pinea* was total when charred crown volume was <50%. No data were found for the more central European species.

4.2.3 Fire frequency

Fire-resistant species are adapted to relatively short fire frequency intervals. This leads to a low accumulation of fuel and thus to a lower intensity and severity. Dendrochronological and paleobotanical studies can help determine the appropriate fire frequency for the persistence of some of the selected fire-resistant species. For example, a relict *P. nigra* forest in the Sierra Turmell in Castellón (eastern Spain) has survived numerous wildfires over several centuries (11 fire dates in the last 172 years) (Fulé et al., 2008). In a *P. sylvestris* forest in Poland and Bielorus, there is evidence of the occurrence low-intensity surface fires at a mean fire interval of 9 ± 7.8 years during the period 1645–2010 (Zin et al., 2015). Paleobotanical studies using sedimentary plant macroremains from two subalpine lakes revealed that to long-term persistence of *P. cembra* was compromised at fire recurrence intervals below 150 years, because this avoids the trees to have enough cambium insulation. Genries determined that fire can affect the cembra pine ecosystem in the Alps if fire frequency is reduced to intervals of less than 80 years. In contrast, recurrence interval of fires lower than 150 years seems to favour *L. decidua* when co-habiting with *P. cembra* while *P. mugo*, abundance seems to be rather stable.

4.2.4 Fire season

For fire-resistant species, we did not find any study that examined post-fire regeneration after crown fires in different seasons. In the case of low to moderate intensity fires, *P. nigra* and *P. sylvestris* are more susceptible to spring than to autumn fires but no data are available on summer conditions (Valor et al., 2017).

4.3 Fire sensitive species and fire regime factors

4.3.1 Pre-fire vegetation: species traits and individual, stand and landscape characteristics.

Fire regimes suitable for forests dominated by fire-sensitive species are characterized by high intensity and infrequent forest fires (Pausas, Bradstock, et al., 2004). The selected fire-sensitive pine species (*P. halepensis* and *P. pinaster*) are characterized by a dual reproductive strategy, producing both serotinous cones and non-serotinous cones. The seeds in serotinous cones remain sealed in the canopy until the rise in temperature dissolves the resin that keeps them sealed and expels the seeds. The seeds then fall to the ground and germinate the following autumn or spring (Daskalidou & Thanos, 1996). The degree of serotiny varies between the selected species and between and within populations of the same species. In Spain, for example, *P. pinaster* has a lower proportion of serotinous cones than *P. halepensis*. The degree of serotiny of *P. halepensis* is more homogeneous in the Iberian Peninsula, while *P. pinaster* shows greater variability along its geographical range (Tapias et al., 2004). *C. vulgaris* can regenerate either from vegetative growth (i.e., sprouting from the root system) or from seed germination from the soil seed bank.

Differences between individuals or populations of the same species are related to the age of individuals and the frequency of crown fires in a given area (Hernández-Serrano et al., 2013). Serotiny is a heritable trait. It is more pronounced in populations with more frequent crown fires than in areas with lower frequency (Hernández-Serrano et al., 2013). At the individual level, the proportion of serotinous cones is higher in young individuals of both species than in adults (Goubitz et al., 2004). For example, *P. halepensis* was found to have 95% serotinous content in young trees and 48% in adult trees in Greece (Daskalidou & Thanos, 1996). The early age of sexual reproduction of both *P. halepensis* and *P. pinaster* and the fact that young trees contain more serotinous cones than adult trees are important for the early accumulation of a large seed bank in the canopy.

The characteristics of the pre-fire stand, such as canopy cover or stand basal area, as well as land-use legacies, can affect the regeneration abundance of fire-sensitive species after a fire. Research has shown that for *P. halepensis*, there is a strong relationship between pre- and post-fire density of pine seedlings, likely due to a greater number of adult pines contributing to the seed bank (Broncano & Retana, 2004; Pausas, et al., 2004). However, no significant effect of land-use legacies has been found (Puerta-Piñero et al., 2012). In contrast, the current post-fire density of *P. pinaster* was found to be largely unpredictable from previous stand history and structure variables in a study developed by Torres et al., (2016). As with fire-resistant species, the recovery of vegetation to prevent erosion depends in the short term on the composition of the understory. A dominant understory of resprouts is likely to recover more quickly than an understory dominated by obligate seeders.

4.3.2 Fire severity

In fire-sensitive species, the degree of fire severity in the canopy is one of the factors determining seed viability for post-fire recruitment. Although *P. halepensis* and *P. pinaster* have reproductive strategies that allow them to recover from fires (serotinous cones), they also have resistance traits such as thick bark (especially *P. pinaster*). This allows them to survive low intensity fires (Fernandes et al., 2008).

Low to moderate fire severity

Mortality from cambium injury is unlikely in *Pinus pinaster* (Fernandes et al., 2008). Based on prescribed burning experiments, the threshold size given by Fernandes et al., (2008) for this species is in the range of 5–10 cm DBH. In the case of *P. halepensis*, there are not many studies on its degree of resistance, but an age of 30 years has been indicated as the threshold to limit cambium injury during prescribed burning experiments (Liacos, 2015). The bark of *P. pinaster* is thicker than the bark of *P. halepensis* Fernandes et al., (2008). Interestingly, *P. halepensis* bark is thicker than *P. sylvestris*, which has been categorized as a fire-resistant species.

High fire severity

In recent decades, the response of *P. halepensis* to fire has been studied in detail. In terms of fire severity, some studies have found no effect on pine seedling abundance (Pausas et al., 2002), while others have reported greater pine regeneration after moderate (Vega et al., 2018) and high severity fires (Elvira et al., 2021; Moya et al., 2020). The differences in the studies can be explained by the different methods used to estimate fire severity. For example, in Pausas et al., (2002), the degree of pine crown consumption was used as an indicator of fire severity: low severity was considered when the crown of trees had > 20% green leaves and high severity when > 80% of crown needles were consumed. The lack of correlation between the density of the seedlings and the severity of the fire could be related to the exposure to heat required to open the serotinous cones that is reached even in low severity fires. In other studies, fire severity was estimated using the delta-normalised burn ratio (dNBR). In all these cases, post-fire regeneration of *P. halepensis* was enhanced by fire severity. In the case of *P. pinaster*, there are fewer studies dealing with recruitment after fire. Martínez et al., (2002) showed that post-fire regeneration was significantly higher in plots affected by surface fire than in plots affected by crown fire. Similarly, Vega et al. (2008) reported that crown damage was negatively (but weakly) related with all parameters of seedling establishment. In another study, Vega et al., (2010) found that scorched trees released the stored seeds more rapidly and to a higher degree than trees with unaffected crown.

In fire-sensitive species, regeneration after fire mostly depends on the canopy seed bank. The role of the soil seed bank and of unburnt trees is small. Temperatures reached in the soil might be too high for survival of the seeds dispersed before the fire and the ability of these species to disperse their seeds

over long distances is low. Maia et al., (2012) concluded that high fire severity can lead to the combustion of the serotinous cones of *P. pinaster*, resulting in lower seed germination. In contrast, less severe fire (scorched and unburned crowns) contributes to increased germination of the seed bank in the soil. According to Escudero et al., (1999), the probability of germination drops to less than 50% at temperatures above 130° C while the role of exposure time is small. In the case of *P. halepensis*, a temperature around 70° C could be decisive for the failure of germination if the exposure time is more than 10 minutes.

C. vulgaris hardly regenerates when the intensities of fire are high because the seeds are killed by burning the moss and litter (MacDonald et al., 1995). Besides, it has also been noted that seed regeneration can be poor where moss and litter layers remain after burning as these provide a poor medium for seedling establishment (Davies, 2006).

4.3.3 Fire frequency

The frequency of high severity fires is a crucial factor for fire-sensitive species, as these species need a sufficient time interval between fires to replenish the soil or the canopy seed bank. In *P. halepensis*, for example, some individuals can form cones as early as 4-8 years of age whereas *P. pinaster* can reach maturity and start producing cones at the age of 4-10 years (Tapias et al., 2004). However, most of trees does not start producing viable cones until 10-20 years after the fire (Tapias et al., 2001; Eugenio & Lloret, 2006). Thus, if the interval between fires is less than 10 years, local extinction of these species may occur. In this sense, the results of Eugenio et al., (2006) show a greater regeneration of *P. halepensis* in places with a lower frequency of fires compared to areas with a higher frequency of fires (25,000 vs. 14,000 seedlings ha⁻¹) and similar results have been obtained by Santana et al., (2010). Recently, Fernández-García et al., (2019) studied the role of fire frequency and fire severity in the recruitment of both species after fire under different environmental conditions. In both species, seedling recruitment was compromised in the driest study area (0.01 seedlings m⁻²), resulting in low seedling cover (0.01%). In *C. vulgaris*, it is estimated that seeds can remain viable in the soil for up to 30-40 years (Gimingham, 1972).

4.3.4 Fire season

Post-fire recruitment for obligate seeders species seems to be reduced by fires outside their historical temporal range (Tangney et al., 2022). Spring fires seems more harmful to seeders species than summer or autumn fires. Short-lived seeders are killed by spring fires before they can produce seed, while long-lived seeds are not yet ready for dispersal in spring (Buhk et al., 2007). Moreover, in case of effective dispersal after a spring fire the seeds have to survive unprotected the upcoming summer and therefore have less chance of germinating in the following rainy season (Chamorro et al., 2013). Tsafrir et al., (2019), for example, found effects of the fire season only in obligate seed plants, with seed survival being lower after spring than after autumn fires.

4.4 Fire tolerant species and fire regime factors

4.4.1 Pre fire vegetation: species traits and individual, stand and landscape characteristics.

Resprouting species can form new shoots after the fire either from below-ground or above-ground structures (Bond & Midgley, 2001).

Above-ground buds can be protected by thick bark or by sinking the buds into the stems (Pausas, 2015). Among the selected species, the two species presenting above-ground buds are *P. canariensis* and *Q. suber*. For *Q. suber* and *P. canariensis*, the size of the tree or the thickness of the bark determines the degree of protection of the epicormic buds. *Q. suber* with a DBH below 12 cm is highly likely to die (Pausas, 1997). For *P. canariensis*, we found only one study in which all examined individuals with a DBH greater than 5 cm resprouted after a fire (Otto et al., 2010).

Belowground buds are buried in the soil, which protects them from heat. They can be stored in roots, root crowns, rhizomes, woody burls, fleshy swellings and belowground caudexes (Pausas et al., 2018). For oaks (*Q. ilex*, *Q. robur* and *Q. pubescens*), *P. tremula* and *B. pendula* there is much evidence of vigorous resprouting after fire through root and stump suckers. The other selected species (*F. sylvatica*, *C. sativa* and *F. angustifolia*) are less well studied, but the data on BROT 2.0 prove they can also resprout after fire. Therefore, we have classified them as fire-tolerant species, even though their ability to resprout may be low. Based on studies of tree mortality after fires found in the review by Espelta et al., (2012), the results of Catry et al., (2010) and the database BROT, we ranked the selected species from lowest to highest probability of dying after fires as follows: *Q. ilex*, *Q. coccifera*, *Q. robur*, *Q. pubescens*, *P. tremula*, *B. pendula*, *F. angustifolia*, *C. sativa* and *F. sylvatica*.

Large specimens usually have a higher survival rate than smaller ones because of the higher number of active buds and the amount of stored carbohydrates (Catry et al., 2010). But specific thresholds for all species have not been found in the literature. In the case of *F. sylvatica*, a DBH of less than 12 cm seems to be critical (Maringer, Ascoli, et al., 2016). For *P. tremula* and *B. pendula*, 100% stem mortality after fire can be expected in individuals with diameters below 6 cm (Ascoli & Bovio, 2010; Brown & Debyle, 2011). For *C. vulgaris*, burning heather at an age between 10 and 15 years is known to allow maximum regeneration of young shoots (Davies & Legg, 2008). Finally, the growth of new shoots in resprouting species is usually positively related to the size of individuals before burning (Espelta et al., 2003).

Post-fire resprouting is also influenced by the stand characteristics prior to the fire and land-use history. For example, the abundance of *Quercus* is highly dependent on its pre-fire stem density, with stands having over 200 stems per hectare typically being dominated by *Quercus* after fire (Torres et al., 2016). In relation to land use history, Puerta-Piñero et al., (2012) found that the abundance

of *Quercus* is strongly influenced by the time since the area was abandoned from agriculture.

4.4.2 Fire severity

Fire tolerant species can survive both low and high intensity fires. Generally high severity does not affect post-fire regeneration (Lloret & López-Soria, 1993). Some studies have shown that more intense forest fires can reduce the ability to resprout, as they can physically damage part of the bud reserve (Casals et al., 2018). Recent studies on *F. sylvatica* have shown that high fire severity, characterised by crown loss above 50% and basal area killed above 60% can result in high mortality, except for polycormic trees (Maringer, Ascoli, et al., 2016).

In the case of *C. vulgaris*, resprouting does not seem to be influenced by the severity or intensity of the fire, as the developmental phase and the loss of *Calluna's* ability to regenerate vegetatively with age are of greater importance (Davies et al., 2010).

4.4.3 Fire frequency

The frequency of fires is an important factor for fire-tolerant species, as these species need a sufficient period between fires to replenish their carbohydrate reserves. In the case of *Q. suber*, a recurrence of about 12 years can impact its post-fire response (Curt et al., 2010). The BROLLA model predicted a decrease from 70% to less than 5% of the total basal area in other *Quercus* species when the fire frequency moves from actual to 5 years (Pausas & Vallejo, 1999). However, in the case of *Q. ilex*, two successive fires in a four-year interval did not limit its ability to resprout (Bonfil et al., 2004). In *Q. robur*, high fire frequency seems to inhibit regeneration, but no specific data on fire recurrence was provided in Monteiro-Henriques & Fernandes (2018). For *F. sylvatica*, basal resprouting from saplings was abundant at fire intervals of 15 years (Moris et al., 2022) whereas in Sweden there is evidence of a *P. tremula* persistence at fire intervals of 30 years (Esseen et al., 1997). No data were found for *P. canariensis*, *Q. pubescens*, *F. angustifolia*, *B. pendula* and *C. sativa*.

Shrubs forms such as *Q. coccifera* had lower biomass in stands affected by recurrent fires (3 fires in 16 years) than in stands affected by only one fire, but they still resprouted (Ferran et al., 2005). In *C. vulgaris*, a fire frequency of 10-15 years allows maximum regeneration of young shoots (Davies & Legg, 2008). In a study conducted in Italy, burning once in 3-6 years appears to be the most appropriate frequency (Ascoli & Bovio, 2010).

4.4.4 Fire season

The timing of fire relative to the active growing season may influence carbohydrate storage and thus the vigour and survival of resprouting species. Although there is limited information on specific studies on the selected resprouting species (*Q. ilex* and *C. vulgaris* mainly), most studies seem to agree that fires at the end of summer or in autumn reduce resprouting and vigour compared

to fires in early summer or spring (Bonfil et al., 2004; Casals et al., 2018; Espelta et al., 2012; Knapp et al., 2005). However, in the case of *C. vulgaris*, new growth after fires was more vigorous in autumn than in spring, but in Scotland where the study was conducted the weather is often dry in early spring, which increases fire damage (Miller & Miles, 2011).

4.5 Topography

The slope or aspect can play an important role in determining erosion risk, but also through its effects on the distribution of soil moisture and thus on the recovery rate of vegetation after fire (Pausas et al., 2004). In Mediterranean areas characterised by a harsh climate, the regeneration rate is usually higher on northern slopes due to lower evapotranspiration and higher moisture content. In wetter climatic areas, the difference may not be relevant.

Fire intolerant

We found only one study dealing with the effect of topography on the recovery of fire-intolerant species after a fire. In Styria (Austria), a threshold slope of about 35-40 degrees seemed to determine whether erosion or regeneration of *P. abies* prevailed (Malowerschnig & Sass, 2014). They conclude that on gentler slopes, grassland was gradually replaced by immature and mature forest, while in steeper areas (approx. > 35-40°) it was degraded to rocks and debris or grassland. No data was found for the effects of aspect.

Fire resistant

In *P. nigra*, south-facing areas seems to reduce post-fire regeneration (Martín-Alcón & Coll, 2016; Martínez-García et al., 2018), specially under steep slopes (Martín-Alcón & Coll, 2016) as also observed in Greece (Christopoulou et al., 2019). Martínez-García et al., (2018) found that slope-aspect did not have an important effect in both seedling emergence and seed predation but it was an important factor for the survival and growth of the emerged pine seedlings. In the case of *P. sylvestris*, a study carried out in Poland after a forest fire showed that regeneration after the fire was stronger on north-facing slopes than on south-facing slopes in the inland dune areas of Central Europe (Sewerniak, 2016). For *L. decidua*, the study of Malowerschnig & Sass, (2014) reported a threshold slope of about 35-40 degrees. In the case of *L. decidua*, a positive relationship between slope and regeneration was found (with high regeneration in slopes > 50%) (Moris et al., 2017). No data was found for *P. pinea*, *P. mugo* and *P. cembra*.

Fire sensitive

The density and height of *P. pinaster* seedlings was higher on north-facing slopes than on south-facing slopes in the three years following a forest fire in north-western Spain (Calvo et al., 2008). This was related to the higher soil moisture found on northern slopes, that is necessary for seedling germination and viability. Similarly, regeneration of *P. halepensis* after fire is greater on north-facing

slopes (Pausas et al., 1999; Pausas, et al., 2004) and on flat/terraced or moderate slopes (Garcia-Jimenez et al., 2017; Pausas et al., 1999). The threshold for slope was set at > 25% in two multicriteria studies (Arianoutsou et al., 2011; Ruiz-Gallardo et al., 2004), but a study developed in Greece show good recovery for slopes between 0-50% (Tsitsoni, 1997). No studies were found for *C. vulgaris*.

Fire tolerant

Studies conducted in Portugal showed that *Q. suber* regeneration was negatively affected by slope. Flat areas were associated with less regeneration (Monteiro-Henriques & Fernandes, 2018) while S, SW and E slopes showed lower crown regeneration than W, NW and NE slopes. The xerophytic character of this species may have led to higher physiological stress and thus higher susceptibility to fire (Catry et al., 2009). Similarly, *Q. robur*, sapling density was lower in southern aspects than in less sun-exposed ones (Monteiro-Henriques & Fernandes, 2018). Studies in Spain on *Q. coccifera* show that resprouting after fire is higher in northern orientations (Pausas & Vallejo, 1999), but in Greece growth is higher in eastern and southern orientations (Konstantinidis et al., 2005). Also in Switzerland, beech regeneration after fire is higher on north-east rather than south-west facing slopes (Maringer, et al., 2016). The resprouting probability of this species did also decrease with slope (Moris et al., 2022). For the other species selected we found no studies.

4.6 Soil characteristics

The recovery of plant species following a fire can be significantly influenced by pre-fire soil properties. However, there has been relatively little investigation into how pre-fire soil properties, such as soil type, bedrock type, soil depth, organic matter content, pH, texture, and nutrient levels, affect post-fire regeneration.

Studies have shown that certain pre-fire soil properties can support faster plant growth and lead to better post-fire recovery. For instance, soils with higher levels of organic matter or nitrogen content, as well as clay soils that can retain more water, tend to promote faster regeneration. In Portugal, recent research has found that vegetation recovery (without distinguishing between species) was most effective in humic cambisols and in soils with low pH (Meneses, 2021). Bedrock type, which is related to soil type, can also have a significant impact on post-fire regeneration. In the Baixo Tamega region of Portugal, for example, shrub formations on granite soils tend to harbor more species and develop more vigorously after fire than those on schist-derived soils (Torres et al., 2017). In Greece, *P. nigra* was more abundant in areas underlain by softer basement rock (Christopoulou et al., 2019). Additionally, Pausas et al., (1999) found that the rate of vegetation regeneration after fire, particularly for *Q. coccifera*, depends on bedrock type. Recovery of vegetation also appears to be greater on soils formed over limestone than on soils over marl. Moreover, López-Soria & Castell (1992) found that resprouting was more common among individuals in deep soils than in

shallow ones. Further research in this area could help inform post-fire management strategies and promote more effective restoration efforts.

4.7 Pre-fire and post-fire climatic conditions

Fire intolerant and resistant species

Research on pre-fire and post-fire climate conditions is limited, with most studies focused on conditions after the fire. For example, in the case of fire intolerant and fire-resistant species such as *P. sylvestris* and *P. abies* in Central Alps, post-fire drought seems to be a critical limiting factor for tree recruitment after fire disturbance (Moser et al., 2010). For both species, the study highlights that the window of opportunity for seedling establishment was short (1-2 years), and moisture deficit was the primary limiting factor for tree recruitment at lower altitudes. The authors found that favourable moisture conditions for seedling establishment in spring were rare at low and medium altitudes. For *L. decidua*, recruitment after fire decreases with the thermal compensated continentality index (Icc), with lower recruitment at Icc values above 27 (Moris et al., 2017). In the case of *P. nigra*, several studies have reported a positive relationship between short-term regeneration and above-average precipitation in the months after fire (Díaz-Delgado et al., 2002, 2003; Viana-Soto et al., 2017).

Fire sensitive species

Regarding fire-sensitive species, most data is limited to *P. halepensis*. A positive relationship between short-term regeneration and above-average precipitation in the months after fire has been found in several studies (Pausas, Ribeiro, et al., 2004; Röder et al., 2008; Viana-Soto et al., 2017). However, a recent study conducted by Elvira et al., (2021) in Catalonia showed that warm conditions before the fire are positively related to pine regeneration, while warm temperatures after the fire have a negative effect on the density of pine trees. Wet conditions after the fire enhanced pine regeneration, while precipitation before the fire had no significant effect. In the case of *P. pinaster*, initially high seedling density decreases significantly after the first summer due to severe drought stress (Calvo et al., 2008).

Fire tolerant species

The resprouting vigour of fire-tolerant species seems to be favoured by precipitation after the fire (Espelta et al., 2012). For instance, in *Q. coccifera*, the identified rainfall episodes after the fire as the decisive positive factor determining resprout development (Konstantinidis et al., 2005). For *F. sylvatica*, spring-summer moisture showed a weak positive effect on beech recruitment (Maringer et al., 2020). Regarding *Q. suber*, post-fire delayed mortality was related to repetitive summer droughts (Curt et al., 2010).

4.8 Other factors

Factors such as competition from pioneers, herbivory and insect outbreaks can also have an important influence on post-fire regeneration but are not considered in this report as the focus here is on variables that can be mapped for WP2.

5 Questionnaire results on factors driving post-fire dynamics

This section presents the results of the questionnaire related to post-fire dynamics (question 9) (Appendix: 8.3 Questionnaire). A total of 35 responses were collected from which 62.8% (n=22) were from the Mediterranean region, 25.7% (n=9) from the continental region, 5.7% (n=2) from the Alpine region, 2.8% (n=1) from the Atlantic region, and 2.8% (n=1) from the temperate region. Most respondents were experts in "fuel management" and "post-fire management" (27.1% and 23.7%, respectively), while 20.3% and 18.6% of respondents were experts in fire behaviour and fire ecology, respectively. The rest of the respondents (10.1%) were experts in other areas of wildland fire research (e.g., risk assessment, social dimension). A large proportion of interviewees were academics (54.2%), followed by fire responders (20.0%), forest managers (17.1%), and other positions (8.5%). The mean years of work experience was 16 years and ranged from 1 to 39 years.

Post-fire dynamics (question 9)

This section presents the results of the questionnaire related with post-fire dynamics (question 9) (Appendix: 8.3 Questionnaire).

Question 9 asked the experts to select a maximum of two vegetation types from a list and then to give a rating from 1 to 10 according to their importance in limiting post-fire recovery in terms of fire regime, pre-fire vegetation, short-term post-fire competition, and climatic, soil and topographic factors. 13 out of 35 experts selected one or two vegetation types. Of all the vegetation types listed, the following were selected: hemiboreal and continental *P. sylvestris* pine forests (n=6), alpine *P. sylvestris* or *P. nigra* in the Alps or *P. uncinata* in the Pyrenees (n=5), *Picea abies* forests (n=3), tall deciduous oak (*Quercus* sp.) forests (n=2), beech (*Fagus* sp.) forests (n=1), *Abies alba* forests (n=1) and mixed *Quercus* sp. and *Fraxinus* forests (n=1). The following forest types were not selected: Hemiboreal mountain pine forests (*Pinus mugo*), subalpine larch forests (*Larix* sp.) and stone pine forests (*Pinus cembra*).

In the case of hemiboreal and continental *P. sylvestris* pine forests, the first five factors, from highest to lowest score, limiting post-fire recovery were fire severity, followed by young forest and high erosion, spring fire and pre-fire drought (Figure 3).




Figure 2. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of hemi boreal and continental Scots pine (*P. sylvestris*) forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.

For alpine *P. sylvestris* or *P. nigra* in the Alps or *P. uncinata* in the Pyrenees, within the first five factors, from highest to lowest score, the first two factors limiting post-fire recovery were related with fire regime factors: fire severity and frequency, followed by post-fire drought. Then, high erosion and torrential rain were punctuated as fire severity, followed by young forest and high erosion, spring fire and pre-fire drought (Figure 3).




Figure 3. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of alpine *P. sylvestris* or *P. nigra* in the Alps or *P. uncinata* in the Pyrenees. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.

For spruce *P. abies*, three of the five top factors were related with climatic conditions (pre- and post-fire drought and torrential rain). Fire frequency was considered the main factor and the age of the forest was positioned in the third place (Figure 4).




Figure 4. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of spruce (*Picea abies*). The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.

In the case of *Quercus* sp. the first five factors affecting post-fire recovery were: post-fire drought, pH acidic, fire severity, anoxy soil and tree vigor and health (Figure 5). It is important to mention that only one or two answers were provided.




Figure 5. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of tall deciduous oak (*Quercus sp.*) forest. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.

For mixed forests of *Quercus sp.* and *Fraxinus* as well as *A. alba*, there was only one answer, which was found in Figure 6 and Figure 7.




Figure 6. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of mixed *Quercus* sp. and *Fraxinus* forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.




Figure 7. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of fir *Abies alba* forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.




Figure 8. Mean importance and standard deviation of punctuation assigned by experts for fire regime, pre-fire vegetation, post-fire short-term competition, climatic, soil and topographic factors influencing post-fire recovery of beech (*Fagus sp.*) forests. The order of the factors was ranked from most important to least important. The number of responses given to each factor is indicated within the bars.

6 Summary: criteria to identify vulnerable areas for restoration before and after fire occurrence

Based on the literature review and the results of the survey presented in section 5, a summary table has been developed outlining key factors, metrics, and thresholds for predicting the areas that may show (at the short-term) slow post-fire plant recovery rates. Restoration action to avoid soil degradation should be prioritized in these areas.

Table 3. Summary of key factors, metrics and thresholds for identifying high priority areas for post-fire restoration. The threshold values given are from the literature review, (-) denotes thresholds not found. The priority of forest types and species was determined based on their fire-related traits and assuming that they burn outside their reference fire regime.

Forest fire type priority	Species priority	Pre-fire factors and associated thresholds						Post and fire-regime factors and associated thresholds		
		Individual characteristics	Pre-fire stand or landscape characteristics	Aspect	Slope	Soil	Pre-fire drought	Fire frequency (years)	Fire season	Post-fire drought
1. Fire intolerant	1. <i>P. abies</i>	DBH <20 cm	Canopy cover ²	South	>35°	Depth ²	Important ²	<200	Summer ¹	Moisture deficit in spring
	2. <i>A. alba</i>	DBH <20 cm	-	-	-	-	-	<200	Summer ¹	-
2. Fire resistant	1. <i>P. mugo</i>	-	-	-	-	-	-	No effect	-	-
	2. <i>L. decidua</i>	-	-	-	-	-	-	>150	-	-
	3. <i>P. cembra</i>	-	-	-	-	-	-	<80	-	-
	4. <i>P. sylvestris</i>	DBH <20 cm	Canopy cover ²	-	-	-	Important ²	>10	Spring	Moisture deficit in spring
	5. <i>P. nigra</i>	DBH < 20 cm	Canopy cover < 50%	South	-	-	-	>10	-	Below-average precipitation in the months after fire
	6. <i>P. pinea</i>	Bark thick. < 2cm	-	-	-	-	-	-	-	-
3. Fire sensitive	1. <i>P. pinaster</i>	Age < 20 years	Undestory of obligate seeders	South	-	-	-	<10	Spring	Drought stress in the first post-fire summer
	2. <i>P.</i>	Age < 20	Low tree	South	>25%	-	No effect	<10	Spring	Warm

	<i>halepensis</i>	years	density Undestory of obligate seeders							conditions
	3. <i>C. vulgaris</i>	Age > 40 years	-	-	-	-	-	> 40	-	-
4. Fire tolerant	1. <i>F. sylvatica</i>	DBH < 12 cm	-	South	Negative effect	-	-	< 15	No data for all species. For resprouting species, the end of summer or autumn seems worse.	-
	2. <i>C. sativa</i>	No specific data but the smaller the individual the less resprouting vigour.	-	-	-	-	-	-		-
	3. <i>F. angustifolia</i>		-	-	-	-	-	-		Continentality index >27
	4. <i>B. pendula</i>		-	-	-	-	-	-		-
	5. <i>P. tremula</i>		-	-	-	-	-	-		-
	6. <i>Q. pubescens</i>		Tree density <200 tree ha ⁻¹	-	-	-	-	-		-
	7. <i>Q. robur</i>		South	-	-	-	-	-		-
	8. <i>Q. coccifera</i>		Time since cropland abandonment	South	-	Bedrock: marls	-	<5		-
	9. <i>Q. ilex</i>		-	-	-	-	<5	-		
	10. <i>C. vulgaris</i>		Age < or > 15 years	-	-	-	-	-	< or > 15	Spring
	11. <i>Q. suber</i>	DBH < 12 cm	-	South, Southwest, East	Flat	-	-	< 12	No data but end summer or fall.	-
	12. <i>P. canariensis</i>	No effect	-	-	-	-	-	-	No data but end summer or fall.	-

¹As compared with winter fires.

²From the questionnaire results.

7 References

- Ananyev, V. A., Timofeeva, V. V., Kryshen', A. M., Pekkoev, A. N., Kostina, E. E., Ruokolainen, A. V., Moshnikov, S. A., Medvedeva, M. V., Polevoi, A. V., & Humala, A. E. (2022). Fire Severity Controls Successional Pathways in a Fire-Affected Spruce Forest in Eastern Fennoscandia. *Forests*, 13(11). <https://doi.org/10.3390/f13111775>
- Arianoutsou, M., Koukoulas, S., & Kazanis, D. (2011). Evaluating Post-Fire Forest Resilience Using GIS and Multi-Criteria Analysis: An Example from Cape Sounion National Park, Greece. *ENVIRONMENTAL MANAGEMENT*, 47(3), 384-397. <https://doi.org/10.1007/s00267-011-9614-7>
- Ascoli, D., & Bovio, G. (2010). Tree encroachment dynamics in heathlands of north-west Italy: the fire regime hypothesis. *iForest - Biogeosciences and Forestry*, 3(5), 137. <https://doi.org/10.3832/IFOR0548-003>
- Bär, A., & Mayr, S. (2020). Bark insulation: Ten Central Alpine tree species compared. *Forest Ecology and Management*, 474(July), 118361. <https://doi.org/10.1016/j.foreco.2020.118361>
- Bär, A., Schröter, D. M., & Mayr, S. (2021). When the heat is on: High temperature resistance of buds from European tree species. *Plant Cell and Environment*, 44(8), 2593-2603. <https://doi.org/10.1111/pce.14097>
- Battipaglia, G., Savi, T., Ascoli, D., Castagneri, D., Esposito, A., Mayr, S., & Nardini, A. (2016). Effects of prescribed burning on ecophysiological, anatomical and stem hydraulic properties in *Pinus pinea* L. . *Tree Physiology*, 36(8), 1019-1031. <https://doi.org/10.1093/treephys/tpw034>
- Bond, W. J., & Midgley, J. J. (2001). Ecology of sprouting in woody plants: the persistence niche. *Trends in ecology & evolution*, 16(1), 45-51.
- Bonfil, C., Cortés, P., Espelta, J. M., & Retana, J. (2004). The role of disturbance in the co-existence of the evergreen *Quercus ilex* and the deciduous *Quercus cerrioides*. *Journal of Vegetation Science*, 15(3), 423-430. <https://doi.org/10.1111/j.1654-1103.2004.tb02280.x>
- Broncano, M. J., & Retana, J. (2004). Topography and forest composition affecting the variability in fire severity and post-fire regeneration occurring after a large fire in the Mediterranean basin. *International Journal of Wildland Fire*, 13(2), 209-216.
- Brown, J. K., & Debyle, N. V. (2011). Fire damage, mortality, and suckering in aspen. <https://doi.org/10.1139/x87-168>, 17(9), 1100-1109. <https://doi.org/10.1139/X87-168>
- Buhk, C., Meyn, A., & Jentsch, A. (2007). The challenge of plant regeneration after fire in the Mediterranean Basin: Scientific gaps in our knowledge on plant strategies and evolution of traits. *Plant Ecology*, 192(1), 1-19.

<https://doi.org/10.1007/s11258-006-9224-2>

- Calvo, L., Santalla, S., Valbuena, L., Marcos, E., Tárrega, R., & Luis-Calabuig, E. (2008). Post-fire natural regeneration of a *Pinus pinaster* forest in NW Spain. *Plant Ecology*, 197(1), 81-90. <https://doi.org/10.1007/s11258-007-9362-1>
- Carter, V. A., Moravcová, A., Chiverrell, R. C., Clear, J. L., Finsinger, W., Dreslerová, D., Halsall, K., & Kuneš, P. (2018). Holocene-scale fire dynamics of central European temperate spruce-beech forests. *Quaternary Science Reviews*, 191, 15-30. <https://doi.org/10.1016/j.quascirev.2018.05.001>
- Casals, P., Valor, T., Rios, A. I., & Shipley, B. (2018). Leaf and bark functional traits predict resprouting strategies of understory woody species after prescribed fires. *Forest Ecology and Management*, 429(July), 158-174. <https://doi.org/10.1016/j.foreco.2018.07.002>
- Catry, F. X., Moreira, F., Duarte, I., & Acácio, V. (2009). Factors affecting post-fire crown regeneration in cork oak (*Quercus suber* L.) trees. *European Journal of Forest Research*, 128(3), 231-240. <https://doi.org/10.1007/s10342-009-0259-5>
- Catry, F. X., Rego, F., Moreira, F., Fernandes, P. M., & Pausas, J. G. (2010). Post-fire tree mortality in mixed forests of central Portugal. *Forest Ecology and Management*, 260(7), 1184-1192. <https://doi.org/10.1016/j.foreco.2010.07.010>
- Chamorro, D., Luna, B., & Moreno, J. M. (2013). Germination response to various temperature regimes of four Mediterranean seeder shrubs across a range of altitudes. *Plant Ecology*, 214(12), 1431-1441. <https://doi.org/10.1007/s11258-013-0264-0>
- Christopoulou, A., Mallinis, G., Vassilakis, E., Farangitakis, G. P., Fyllas, N. M., Kokkoris, G. D., & Arianoutsou, M. (2019). Assessing the impact of different landscape features on post-fire forest recovery with multitemporal remote sensing data: The case of Mount Taygetos (southern Greece). *International Journal of Wildland Fire*, 28(7), 521-532. <https://doi.org/10.1071/WF18153>
- Curt, T., Bertrand, R., Borgniet, L., Ferrieux, T., & Marini, E. (2010). The impact of fire recurrence on populations of *Quercus suber* in southeastern France. *VI International Conference on Forest Fire Research*, 10-p.
- Daskalidou, E. N., & Thanos, C. A. (1996). Aleppo pine (*Pinus halepensis*) postfire regeneration: the role of canopy and soil seed banks. *International Journal of Wildland Fire*, 6(2), 59-66.
- Davies, G. M. (2006). *Fire behaviour and impact on heather moorland*.
- Davies, G. M., & Legg, C. J. (2008). The effect of traditional management burning on lichen diversity. *Applied Vegetation Science*, 11(4), 529-538. <https://doi.org/10.3170/2008-7-18566>
- Díaz-Delgado, R., Lloret, F., & Pons, X. (2003). Influence of fire severity on plant regeneration by means of remote sensing imagery. *International Journal of Remote Sensing*, 24(8), 1751-1763.

- Díaz-Delgado, R., Lloret, F., Pons, X., & Terradas, J. (2002). Satellite evidence of decreasing resilience in Mediterranean plant communities after recurrent wildfires. *Ecology*, *83*(8), 2293-2303.
- Dupire, S., Curt, T., Bigot, S., & Fréjaville, T. (2019). Vulnerability of forest ecosystems to fire in the French Alps. *European Journal of Forest Research*, *138*(5), 813-830. <https://doi.org/10.1007/s10342-019-01206-1>
- Elvira, N. J., Lloret, F., Jaime, L., Margalef-Marrase, J., Pérez Navarro, M. Á., & Batllori, E. (2021). Species climatic niche explains post-fire regeneration of Aleppo pine (*Pinus halepensis* Mill.) under compounded effects of fire and drought in east Spain. *Science of the Total Environment*, *798*, 149308. <https://doi.org/10.1016/j.scitotenv.2021.149308>
- Espelta, J. M., Barbati, A., Quevedo, L., Tarrega, R., Navascues, P., Bonfil, C., Peguero, G., Fernandez-Martinez, M., & Rodrigo, A. (2012). Post-Fire Management of Mediterranean Broadleaved Forests. En F. Moreira, M. Arianoutsou, P. Corona, & J. D. L. Heras (Eds.), *POST-FIRE MANAGEMENT AND RESTORATION OF SOUTHERN EUROPEAN FORESTS* (Vol. 24, pp. 171-194). https://doi.org/10.1007/978-94-007-2208-8_8
- Espelta, J. M., Retana, J., & Habrouk, A. (2003). Resprouting patterns after fire and response to stool cleaning of two coexisting Mediterranean oaks with contrasting leaf habits on two different sites. *Forest Ecology and Management*, *179*(1-3), 401-414.
- Esseen, P.-A., Ehnström, B., Ericson, L., & Sjöberg, K. (1997). Boreal forests. *Ecological bulletins*, 16-47.
- Fernandes, P. M., Vega, A., & Jime, E. (2008). *Forest Ecology and Management Fire resistance of European pines*. 1-10. <https://doi.org/10.1016/j.foreco.2008.04.032>
- Fernández-García, V., Fulé, P. Z., Marcos, E., & Calvo, L. (2019). The role of fire frequency and severity on the regeneration of Mediterranean serotinous pines under different environmental conditions. *Forest Ecology and Management*, *444*(March), 59-68. <https://doi.org/10.1016/j.foreco.2019.04.040>
- Ferran, A., Delitti, W., & Vallejo, V. R. (2005). Effects of fire recurrence in *Quercus coccifera* L. shrublands of the Valencia Region (Spain): II. plant and soil nutrients. *Plant Ecology*, *177*, 71-83.
- Feurdean, A., Florescu, G., Vannièrè, B., Tanțău, I., O'Hara, R. B., Pfeiffer, M., Hutchinson, S. M., Gałka, M., Moskal-del Hoyo, M., & Hickler, T. (2017). Fire has been an important driver of forest dynamics in the Carpathian Mountains during the Holocene. *Forest Ecology and Management*, *389*, 15-26. <https://doi.org/10.1016/j.foreco.2016.11.046>
- Feurdean, A., Tonkov, S., Pfeiffer, M., Panait, A., Warren, D., Vannièrè, B., & Marinova, E. (2019). Fire frequency and intensity associated with functional traits of dominant forest type in the Balkans during the Holocene. *European Journal of Forest Research*, *138*(6), 1049-1066. <https://doi.org/10.1007/s10342->

019-01223-0

- Finsinger, W., Fevre, J., Orbán, I., Pál, I., Vincze, I., Hubay, K., Birks, H. H., Braun, M., Tóth, M., & Magyari, E. K. (2018). Holocene fire-regime changes near the treeline in the Retezat Mts. (Southern Carpathians, Romania). *Quaternary International*, 477(May), 94-105. <https://doi.org/10.1016/j.quaint.2016.04.029>
- Frejaville, T., Curt, T., & Carcaillet, C. (2013). Bark flammability as a fire-response trait for subalpine trees. *Frontiers in plant science*, 4, 466.
- Fulé, P. Z., Ribas, M., Gutiérrez, E., Vallejo, R., & Kaye, M. W. (2008). Forest structure and fire history in an old *Pinus nigra* forest, eastern Spain. *Forest Ecology and Management*, 255(3-4), 1234-1242.
- Garcia-Jimenez, R., Palmero-Iniesta, M., & Maria Espelta, J. (2017). Contrasting Effects of Fire Severity on the Regeneration of *Pinus halepensis* Mill. and Resprouter Species in Recently Thinned Thickets. *FORESTS*, 8(3). <https://doi.org/10.3390/f8030055>
- Genries, A., Mercier, L., Lavoie, M., Muller, S. D., Radakovitch, O., & Carcaillet, C. (2009). The effect of fire frequency on local cembra pine populations. *Ecology*, 90(2), 476-486. <https://doi.org/10.1890/07-1740.1>
- Gimingham, C. H. (1972). *Ecology of heathlands*.
- Goubitz, S., Nathan, R., Roitemberg, R., Shmida, A., & Ne'eman, G. (2004). Canopy seed bank structure in relation to: fire, tree size and density. *Plant Ecology*, 173, 191-201.
- Hernández-Serrano, A., Verdú, M., González-Martínez, S. C., & Pausas, J. G. (2013). Fire structures pine Serotiny at different scales. *American Journal of Botany*, 100(12), 2349-2356. <https://doi.org/10.3732/ajb.1300182>
- Knapp, E. E., Keeley, J. E., Ballenger, E. A., & Brennan, T. J. (2005). Fuel reduction and coarse woody debris dynamics with early season and late season prescribed fire in a Sierra Nevada mixed conifer forest. *Forest Ecology and Management*, 208(1-3), 383-397.
- Konstantinidis, P., Tsiourlis, G., & Galatsidas, S. (2005). Effects of wildfire season on the resprouting of kermes oak (*Quercus coccifera* L.). *Forest Ecology and Management*, 208(1-3), 15-27. <https://doi.org/10.1016/j.foreco.2004.09.021>
- Leys, B., Carcaillet, C., Blarquez, O., Lami, A., Musazzi, S., & Trevisan, R. (2014). Resistance of mixed subalpine forest to fire frequency changes: The ecological function of dwarf pine (*Pinus mugo* ssp. *mugo*). *Quaternary Science Reviews*, 90, 60-68. <https://doi.org/10.1016/j.quascirev.2014.02.023>
- Liacos, L. G. (2015). Present Studies and History of Burning in Greece. *Fire Ecology*, 11(3), 3-13. <https://doi.org/10.1007/BF03400631>
- Lloret, F., & López-Soria, L. (1993). Resprouting of *Erica multiflora* after experimental fire treatments. *Journal of Vegetation Science*, 4(3), 367-374.

- López-Soria, L., & Castell, C. (1992). Comparative genet survival after fire in woody Mediterranean species. *Oecologia*, 91(4), 493-499. <https://doi.org/10.1007/BF00650321>
- MacDonald, A. J., Kirkpatrick, A. H., Hester, A. J., & Sydes, C. (1995). Regeneration by Natural Layering of Heather (*Calluna vulgaris*): Frequency and Characteristics in Upland Britain. *Journal of Applied Ecology*, 32(1), 85-99. <https://doi.org/10.2307/2404418>
- Madrigal, J., Souto-García, J., Calama, R., Guijarro, M., Picos, J., & Hernando, C. (2019). Resistance of *Pinus pinea* L. bark to fire. *International Journal of Wildland Fire*, 28(5), 342-353. <https://doi.org/10.1071/WF18118>
- Malowerschnig, B., & Sass, O. (2014). Long-term vegetation development on a wildfire slope in Innerzwain (Styria, Austria). *Journal of Forestry Research*, 25(1), 103-111. <https://doi.org/10.1007/s11676-014-0435-4>
- Maringer, J., Ascoli, D., Kuffer, N., Schmidlein, S., & Conedera, M. (2016). What drives European beech (*Fagus sylvatica* L.) mortality after forest fires of varying severity? *FOREST ECOLOGY AND MANAGEMENT*, 368, 81-93. <https://doi.org/10.1016/j.foreco.2016.03.008>
- Maringer, J., Conedera, M., Ascoli, D., Schmatz, D. R., & Wohlgemuth, T. (2016). Resilience of European beech forests (*Fagus sylvatica* L.) after fire in a global change context. *International Journal of Wildland Fire*, 25(6), 699-710. <https://doi.org/10.1071/WF15127>
- Maringer, J., Wohlgemuth, T., Hacket-Pain, A., Ascoli, D., Berretti, R., & Conedera, M. (2020). Drivers of persistent post-fire recruitment in European beech forests. *SCIENCE OF THE TOTAL ENVIRONMENT*, 699. <https://doi.org/10.1016/j.scitotenv.2019.134006>
- Martin-Alcon, S., & Coll, L. (2016). Unraveling the relative importance of factors driving post-fire regeneration trajectories in non-serotinous *Pinus nigra* forests. *FOREST ECOLOGY AND MANAGEMENT*, 361, 13-22. <https://doi.org/10.1016/j.foreco.2015.11.006>
- Martín-Alcón, S., & Coll, L. (2016). Unraveling the relative importance of factors driving post-fire regeneration trajectories in non-serotinous *Pinus nigra* forests. *Forest Ecology and Management*, 361, 13-22.
- Martinez-Garcia, E., Miettinen, H., Rubio, E., Antonio Garcia-Morote, F., Andres-Abellan, M., & Ramon Lopez-Serrano, F. (2018). Effects of post-fire management practices and slope-aspect on medium-term Spanish black pine regeneration: implications of using a direct seeding strategy in burnt areas. *EUROPEAN JOURNAL OF FOREST RESEARCH*, 137(4), 527-540. <https://doi.org/10.1007/s10342-018-1121-4>
- Martínez, E., Madrigal, J., Hernando, C., Guijarro, M., Vega, J. A., Pérez-Gorostiaga, P., Fonturbel, M. T., Cuiñas, P., Alonso, M., & Beloso, M. C. (2002). Effect of fire intensity on seed dispersal and early regeneration in a *Pinus pinaster* forest.

Forest fire research and wildland fire safety: Proceedings of IV International Conference on Forest Fire Research 2002 Wildland Fire Safety Summit, Luso, Coimbra, Portugal, 18-23 November 2002.

- Matt Davies, G., Adam Smith, A., MacDonald, A. J., Bakker, J. D., & Legg, C. J. (2010). Fire intensity, fire severity and ecosystem response in heathlands: Factors affecting the regeneration of *Calluna vulgaris*. *Journal of Applied Ecology*, 47(2), 356-365. <https://doi.org/10.1111/j.1365-2664.2010.01774.x>
- Meneses, B. M. (2021). Vegetation recovery patterns in burned areas assessed with landsat 8 oli imagery and environmental biophysical data. *Fire*, 4(4). <https://doi.org/10.3390/fire4040076>
- Miller, A. G. R., & Miles, J. (2011). (L .) Hull) At Different Ages and Seasons in North-East Scotland. *Society*, 7(1), 51-60.
- Monteiro-Henriques, T., & Fernandes, P. M. (2018). Regeneration of native forest species in Mainland Portugal: Identifying main drivers. *Forests*, 9(11). <https://doi.org/10.3390/f9110694>
- Moreira, F., Arianoutsou, M., Vallejo, V. R., de las Heras, J., Corona, P., Xanthopoulos, G., Fernandes, P., & Papageorgiou, K. (2012). *Setting the Scene for Post-Fire Management* (pp. 1-19). https://doi.org/10.1007/978-94-007-2208-8_1
- Moris, J. V., Vacchiano, G., Ravetto Enri, S., Lonati, M., Motta, R., & Ascoli, D. (2017a). Resilience of European larch (*Larix decidua* Mill.) forests to wildfires in the western Alps. En *New Forests* (Vol. 48, Número 5). <https://doi.org/10.1007/s11056-017-9591-7>
- Moris, J. V, Berretti, R., Bono, A., Sino, R., Minotta, G., Garbarino, M., Motta, R., Vacchiano, G., Maringer, J., Conedera, M., & Ascoli, D. (2022). Resprouting in European beech confers resilience to high-frequency fire. *FORESTRY*. <https://doi.org/10.1093/forestry/cpac018>
- Moris, J. V, Vacchiano, G., Ravetto Enri, S., Lonati, M., Motta, R., & Ascoli, D. (2017b). Resilience of European larch (*Larix decidua* Mill.) forests to wildfires in the western Alps. *NEW FORESTS*, 48(5), 663-683. <https://doi.org/10.1007/s11056-017-9591-7>
- Moser, B., Temperli, C., Schneiter, G., & Wohlgemuth, T. (2010). Potential shift in tree species composition after interaction of fire and drought in the Central Alps. *European Journal of Forest Research*, 129(4), 625-633. <https://doi.org/10.1007/s10342-010-0363-6>
- Nolan, R. H., Collins, L., Leigh, A., Ooi, M. K. J., Curran, T. J., Fairman, T. A., Resco de Dios, V., & Bradstock, R. (2021). Limits to post-fire vegetation recovery under climate change. *Plant Cell and Environment*, 44(11), 3471-3489. <https://doi.org/10.1111/pce.14176>
- Ordóñez, J. L., Molowny-Horas, R., & Retana, J. (2006). A model of the recruitment

- of *Pinus nigra* from unburned edges after large wildfires. *Ecological Modelling*, 197(3-4), 405-417.
- Ordóñez, J. L., Retana, J., & Espelta, J. M. (2005). Effects of tree size, crown damage, and tree location on post-fire survival and cone production of *Pinus nigra* trees. *Forest Ecology and Management*, 206(1-3), 109-117.
- Otto, R., Garcia-del-Rey, E., Munoz, P. G., & Fernandez-Palacios, J. M. (2010). The effect of fire severity on first-year seedling establishment in a *Pinus canariensis* forest on Tenerife, Canary Islands. *EUROPEAN JOURNAL OF FOREST RESEARCH*, 129(4), 499-508. <https://doi.org/10.1007/s10342-009-0347-6>
- Pausas, J. G. (1997). Resprouting of *Quercus suber* in NE Spain after fire. *Journal of Vegetation Science*, 8(5), 703-706. <https://doi.org/10.2307/3237375>
- Pausas, J. G. (2015). Bark thickness and fire regime. *Functional Ecology*, 29(3), 315-327.
- Pausas, J. G., Bradstock, R. A., Keith, D. A., & Keeley, J. E. (2004). Plant functional traits in relation to fire in crown-fire ecosystems. *Ecology*, 85(4), 1085-1100.
- Pausas, J. G., Carbo, E., Caturla, R. N., Gil, J. M., & Vallejo, R. (1999). Post-fire regeneration patterns in the eastern Iberian Peninsula. *ACTA OECOLOGICA-INTERNATIONAL JOURNAL OF ECOLOGY*, 20(5), 499-508. [https://doi.org/10.1016/S1146-609X\(00\)86617-5](https://doi.org/10.1016/S1146-609X(00)86617-5)
- Pausas, J. G., & Keeley, J. E. (2014). Evolutionary ecology of resprouting and seeding in fire-prone ecosystems. *New Phytologist*, 204(1), 55-65. <https://doi.org/10.1111/nph.12921>
- Pausas, J. G., Lamont, B. B., Paula, S., Appezzato-da-Glória, B., & Fidelis, A. (2018). Unearthing belowground bud banks in fire-prone ecosystems. *New Phytologist*, 217(4), 1435-1448. <https://doi.org/10.1111/nph.14982>
- Pausas, J. G., Ribeiro, E., & Vallejo, R. (2004). Post-fire regeneration variability of *Pinus halepensis* in the eastern Iberian Peninsula. *FOREST ECOLOGY AND MANAGEMENT*, 203(1-3), 251-259. <https://doi.org/10.1016/j.foreco.2004.07.061>
- Pausas, J. G., & Vallejo, V. R. (1999). The role of fire in European Mediterranean ecosystems. *Remote Sensing of Large Wildfires: in the European Mediterranean Basin*, 3-16.
- Pividori, M., Giannetti, F., Barbatì, A., & Chirici, G. (2016). European Forest Types: tree species matrix. *European Atlas of Forest Tree Species*, e01f162.
- Puerta-Piñero, C., Espelta, J. M., Sánchez-Humanes, B., Rodrigo, A., Coll, L., & Brotons, L. (2012). History matters: Previous land use changes determine post-fire vegetation recovery in forested Mediterranean landscapes. *Forest Ecology and Management*, 279, 121-127. <https://doi.org/10.1016/j.foreco.2012.05.020>
- Rigolot, E. (2004). Predicting postfire mortality of *Pinus halepensis* Mill. and *Pinus*

- pinia L. *Plant Ecology*, 171(1-2), 139-151.
- Röder, A., Hill, J., Duguay, B., Alloza, J. A., & Vallejo, R. (2008). Using long time series of Landsat data to monitor fire events and post-fire dynamics and identify driving factors. A case study in the Ayora region (eastern Spain). *Remote Sensing of Environment*, 112(1), 259-273. <https://doi.org/10.1016/j.rse.2007.05.001>
- Rodrigo, A., Quintana, V., & Retana, J. (2007). Fire reduces *Pinus pinia* distribution in the northeastern Iberian Peninsula. *Ecoscience*, 14(1), 23-30. [https://doi.org/10.2980/1195-6860\(2007\)14\[23:FRPPDI\]2.0.CO;2](https://doi.org/10.2980/1195-6860(2007)14[23:FRPPDI]2.0.CO;2)
- Rogers, B. M., Soja, A. J., Goulden, M. L., & Randerson, J. T. (2015). Influence of tree species on continental differences in boreal fires and climate feedbacks. *Nature Geoscience*, 8(3), 228-234. <https://doi.org/10.1038/ngeo2352>
- Ruiz-Gallardo, J. R., Castaño, S., & Calera, A. (2004). Application of remote sensing and GIS to locate priority intervention areas after wildland fires in Mediterranean systems: a case study from south-eastern Spain. *International Journal of Wildland Fire*, 13(3), 241-252.
- Sewerniak, P. (2016). Differences in early dynamics and effects of slope aspect between naturally regenerated and planted *Pinus sylvestris* woodland on inland dunes in Poland. *IForest*, 9(6), 875-882. <https://doi.org/10.3832/ifor1728-009>
- Tangney, R., Paroissien, R., Le Breton, T. D., Thomsen, A., Doyle, C. A. T., Ondik, M., Miller, R. G., Miller, B. P., & Ooi, M. K. J. (2022). Success of post-fire plant recovery strategies varies with shifting fire seasonality. *Communications Earth and Environment*, 3(1), 1-9. <https://doi.org/10.1038/s43247-022-00453-2>
- Tapias, R., Climent, J., Pardos, J. A., & Gil, L. (2004). Life histories of Mediterranean pines. *Plant Ecology*, 171(1-2), 53-68. <https://doi.org/10.1023/B:VEGE.0000029383.72609.f0>
- Tavşanoğlu, Ç., & Pausas, J. G. (2018). Data Descriptor: A functional trait database for Mediterranean Basin plants. *Scientific Data*, 5, 1-18. <https://doi.org/10.1038/sdata.2018.135>
- Tinner, W., Conedera, M., Gobet, E., Hubschmid, P., Wehrli, M., & Ammann, B. (2000). A palaeoecological attempt to classify fire sensitivity of trees in the southern Alps. *Holocene*, 10(5), 565-574. <https://doi.org/10.1191/095968300674242447>
- Torres, I., Pérez, B., Quesada, J., Viedma, O., & Moreno, J. M. (2016). Forest shifts induced by fire and management legacies in a *Pinus pinaster* woodland. *Forest Ecology and Management*, 361(October 2017), 309-317. <https://doi.org/10.1016/j.foreco.2015.11.027>
- Torres, J., Marques, J., Alves, P., Costa, H., & Honrado, J. (2017). Local lithological drivers of post-fire vegetation recovery and implications for fire-prone

- regions. *ECOLOGICAL RESEARCH*, 32(1), 37-49. <https://doi.org/10.1007/s11284-016-1415-2>
- Tsafir, A., Osem, Y., Shemesh, H., Carmel, Y., Soref, C., & Ovadia, O. (2019). Fire season modifies the perennial plant community composition through a differential effect on obligate seeders in eastern Mediterranean woodlands. *Applied Vegetation Science*, 22(1), 115-126. <https://doi.org/10.1111/avsc.12408>
- Tsitsoni, T. (1997). Conditions determining natural regeneration after wildfires in the *Pinus halepensis* (Miller, 1768) forests of Kassandra peninsula (north Greece). *FOREST ECOLOGY AND MANAGEMENT*, 92(1-3), 199-208. [https://doi.org/10.1016/S0378-1127\(96\)03909-6](https://doi.org/10.1016/S0378-1127(96)03909-6)
- Vallejo, V. R., Arianoutsou, M., & Moreira, F. (2012). Post-Fire Management and Restoration of Southern European Forests. *Post-fire forest management in southern Europe: a COST action for gathering and disseminating scientific knowledge*, 24(January), 93-119. <https://doi.org/10.1007/978-94-007-2208-8>
- Valor, T., Coll, L., Pique, M., Dupuy, J. ., & Casals, P. (2023). *FIRE-RES Ecological factors driving resistant and resilient landscapes to high intensity and extreme wildfire events. Deliverable D1.11 FIRE-RES project.*
- Valor, T., González-Olabarria, J. R., Piqué, M., & Casals, P. (2017). The effects of burning season and severity on the mortality over time of *Pinus nigra* spp. *salzmannii* (Dunal) Franco and *P. sylvestris* L. *Forest Ecology and Management*, 406(June), 172-183. <https://doi.org/10.1016/j.foreco.2017.08.027>
- Vega, J. A., Fernández, C., Pérez-Gorostiaga, P., & Fonturbel, T. (2010). Response of maritime pine (*Pinus pinaster* Ait.) recruitment to fire severity and post-fire management in a coastal burned area in Galicia (NW Spain). *Plant Ecology*, 206(2), 297-308. <https://doi.org/10.1007/s11258-009-9643-y>
- Viana-Soto, A., Aguado, I., & Martinez, S. (2017). Assessment of Post-Fire Vegetation Recovery Using Fire Severity and Geographical Data in the Mediterranean Region (Spain). *ENVIRONMENTS*, 4(4). <https://doi.org/10.3390/environments4040090>
- Zin, E., Drobyshev, I., Bernacki, D., & Niklasson, M. (2015). Dendrochronological reconstruction reveals a mixed-intensity fire regime in *Pinus sylvestris*-dominated stands of Białowież'a Forest, Belarus and Poland. *Journal of Vegetation Science*, 26(5), 934-945. <https://doi.org/10.1111/jvs.12290>

8 Appendices

8.1 Preliminary dominant species selected

Table 3. Preliminary selection of species for conducting the literature review on the influence of different factors on post-fire recovery. The selection of species was based on the work of Pividori et al., (2016), who classified the species presence for each European forest type (EFT) of the EFT classification into three classes: the species is abundant and dominant in the EFT; the species presence in the EFT is either secondary or predominant, but under particular and uncharacteristic ecological conditions of the EFT; the presence in the EFT is in some cases both dominant and secondary. Only species that were abundant and dominant in a given EFT were selected from these categories.

Dominat species	Category	Forest type
<i>Abies alba</i>	2. Hemiboreal forest and nemoral coniferous and mixed broadleaved-coniferous forest	2.8 Nemoral silver fir
	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.6 Mediterranean and Anatolian fir forest
<i>Picea abies</i>	1. Boreal forest	1.1 Spruce and spruce-birch boreal forest
	2. Hemiboreal forest and nemoral coniferous and mixed broadleaved-coniferous forest	2.1 Hemiboreal forest
		2.3 Nemoral spruce forest
	3. Alpine coniferous forest	3.2 Subalpine and mountainous spruce and mountainous mixed spruce-silver fir forest
	11. Mire and swamp forest	11.1 Spruce mire forest
<i>Larix decidua</i>	3. Alpine coniferous forest	3.1 Subalpine larch-arolla pine and dwarf pine forest
<i>Pinus pinaster</i>	2. Hemiboreal forest and nemoral coniferous and mixed broadleaved-coniferous forest	2.7 Atlantic maritime pine forest
	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.1.1 Mediterranean pine forest - Pinus pinaster
<i>Pinus nigra</i>	2. Hemiboreal forest and nemoral coniferous and mixed broadleaved-coniferous forest	2.4 Nemoral Black pine forest
	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.2.1 Mediterranean Black pine forest
		10.2.2 Anatolian Black pine forest
<i>Pinus sylvestris</i>	1. Boreal forest	1.2 Pine and pine-birch boreal forest
	2. Hemiboreal forest and nemoral coniferous and mixed broadleaved-coniferous forest	2.1 Hemiboreal forest
		2.2 Nemoral Scots pine forest
		2.5 Mixed Scots pine-birch forest
	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian	10.4 Mediterranean and Anatolian Scots pine forest

	regions	
	11. Mire and swamp forest	11.2 Pine mire forest
<i>Pinus mugo</i>	3. Alpine coniferous forest	3.1 Subalpine larch-arella pine and dwarf pine forest
<i>Pinus halepensis</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.1.2 Mediterranean pine forest - <i>Pinus halepensis</i>
<i>Pinus pinea</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.1.3 Mediterranean pine forest - <i>Pinus pinea</i>
<i>Pinus cembra</i>	3. Alpine coniferous forest	3.1 Subalpine larch-arella pine and dwarf pine forest
<i>Pinus canariensis</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.3 Canarian pine forest
<i>Cupressus sempervirens</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.8 Cypress forest
<i>Cedrus libani</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.9 Cedar forest
<i>Cedrus brevifolia</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.9 Cedar forest
<i>Tetraclinis articulata</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.10 Tetraclinis articulata stands
<i>Taxus baccata</i>	10. Coniferous forests of the Mediterranean, Anatolian and Macaronesian regions	10.11 Mediterranean yew stands
<i>Populus tremula</i>	11. Mire and swamp forest	11.6 Aspen swamp forest
	13. Non riverine alder, birch, or aspen forest	13.4 Aspen forest
<i>Betula pubescens</i>	3. Alpine coniferous forest	3.4 Mountainous birch forest
	13. Non riverine alder, birch, or aspen forest	13.3 Birch forest
<i>Betula pendula</i>	11. Mire and swamp forest	11.4 Birch swamp forest
	13. Non riverine alder, birch, or aspen forest	13.3 Birch forest
<i>Alnus viridis</i>	13. Non riverine alder, birch, or aspen forest	13.1 Alder forest
<i>Alnus cordata</i>	13. Non riverine alder, birch, or aspen forest	13.1 Alder forest
		13.2 Italian alder forest
<i>Carpinus betulus</i>	5. Mesophytic deciduous forest	5.1 Pedunculate oak-hornbeam forest
		5.2 Sessile oak-hornbeam forest
<i>Carpinus orientalis</i>	6. Beech forest	6.2 Atlantic and subatlantic lowland beech forest
		8.8.4 Oriental hornbeam (<i>Carpinus orientalis</i>) forest
<i>Ostrya carpinifolia</i>	8. Thermophilous deciduous forest	8.8.3 Hop-hornbeam (<i>Ostrya carpinifolia</i>) forest
<i>Fagus sylvatica</i>	6. Beech forest	6.1 Lowland beech forest of southern Scandinavia and north central Europe

		6.2 Atlantic and subatlantic lowland beech forest
		6.3 Subatlantic submountainous beech forest
		6.4 Central European submountainous beech forest
		6.5 Carpathian submountainous beech forest
		6.6 Illyrian submountainous beech forest
		6.7 Moesian submountainous beech forest
	7. Mountainous beech forest	7.1 South western European mountainous beech forest
	7. Mountainous beech forest	7.2 Central European mountainous beech forest
	7. Mountainous beech forest	7.3 Apennine-Corsican mountainous beech forest
	7. Mountainous beech forest	7.4 Illyrian mountainous beech forest
	7. Mountainous beech forest	7.5 Carpathian mountainous beech forest
	7. Mountainous beech forest	7.6 Moesian mountainous beech forest
<i>Fagus moesiaca</i>	6. Beech forest	6.5 Carpathian submountainous beech forest
	6. Beech forest	6.7 Moesian submountainous beech forest
	7. Mountainous beech forest	7.4 Illyrian mountainous beech forest
		7.6 Moesian mountainous beech forest
7.7 Crimean mountainous beech forest		
<i>Fagus orientalis</i>	6. Beech forest	6.5 Carpathian submountainous beech forest
<i>Castanea sativa</i>	8. Thermophilous deciduous forest	8.7 Chestnut forest
<i>Quercus coccifera</i>	9. Broadleaved evergreen forest	9.1 Mediterranean evergreen oak forest
<i>Quercus ilex</i>	9. Broadleaved evergreen forest	9.1 Mediterranean evergreen oak forest
<i>Quercus suber</i>	9. Broadleaved evergreen forest	9.1 Mediterranean evergreen oak forest
<i>Quercus trojana</i>	8. Thermophilous deciduous forest	8.5 Macedonian oak forest
<i>Quercus robur</i>	11. Mire and swamp forest	11.5 Pedunculate oak swamp forest
<i>Quercus pubescens</i>	8. Thermophilous deciduous forest	8.1.1 Downy oak forest - western
		8.1.2 Downy oak forest - Italian
		8.1.3 Downy oak forest - Greek, Anatolian
		8.1.4 Downy oak forest - steppe

<i>Ulmus grabra</i>	5. Mesophytic deciduous forest	5.8 Ravine and slope forest
<i>Celtis australis</i>	8. Thermophilous deciduous forest	8.8.7 Celtis australis forest
<i>Acer pseudoplatanus</i>	5. Mesophytic deciduous forest	5.8 Ravine and slope forest
<i>Tilia cordata</i>	5. Mesophytic deciduous forest	5.7 Lime forest
<i>Tilia platyphyllos</i>	5. Mesophytic deciduous forest	5.8 Ravine and slope forest
<i>Fraxinus excelsior</i>	5. Mesophytic deciduous forest	5.8 Ravine and slope forest
<i>Fraxinus angustifolia</i>	8. Thermophilous deciduous forest	8.8.1 Thermophilous ash forest

8.2 Literature review: search strategy

For the species that were abundant and dominant in a given European Forest Type we conducted a systematic literature review and searched in the ISI Web of Knowledge for European studies that addressed how, fire regime, soil characteristics, topography, landscape structure influences post-fire regeneration. The keyword search used various combinations of relevant terms (Table 4).

Table 4. Keyword searches for each fuel-related factors associated with stand and landscape resistance.

Factor	Keywords
Fire severity	Species scientific name AND fire severity AND post-fire regeneration OR post-fire recovery OR resistance
Fire frequency	Species scientific name AND fire frequency AND post-fire regeneration OR post-fire recovery OR resistance
Fire season	Species scientific name AND fire season AND post-fire regeneration OR post-fire recovery OR resistance
Aspect	Species scientific name AND aspect AND post-fire regeneration OR post-fire recovery OR resistance
Slope	Species scientific name AND slope AND post-fire regeneration OR post-fire recovery OR resistance
Climate	Species scientific name AND fire severity AND post-fire regeneration OR post-fire recovery OR resistance

8.3 Questionnaire

FACTORS DRIVING RESISTANT AND RESILIENT LANDSCAPES TO EXTREME WILDFIRE EVENTS AND POST-FIRE DYNAMICS

FIRE-RES -Innovative technologies and socio-ecological-economic solutions for re resilient territories in Europe - ([https:// re-res.eu/](https://re-res.eu/)) is an ongoing **Horizon 2020** project (2021-2025). **FIRE-RES** aims to promote the implementation of an integrated re management approach and support the transition to more resilient landscapes and communities to extreme wild re events (EWE) in Europe. The mission of **FIRE-RES** is to promote the European Union's socio-ecological transition to a resilient continent through the development of a series of innovation actions.

One of the objectives is to develop general recommendations and management alternatives **to promote adaptive management for resilient landscapes to EWE**. To this end, we conducted a systematic literature review to identify ecological factors, metrics, and thresholds that determine landscape and stand resistance and resilience to EWE and post- re dynamics. Most of the identified factors and thresholds were found for southern Mediterranean countries and are related to large and intense wild res only a few for EWE. Therefore, **the main objective of the questionnaire is to collect experts' views on the thresholds of different fuel-related factors to prevent the development of EWE and data for Central and Northern Europe**, where wildfires are not yet so common in these relatively humid regions, but where an increase in the frequency and intensity of wild res can be expected. This is a key issue because the importance of fuel-related factors and thresholds that determine resistance and resilience may vary by biogeographic area and wildfire type. By examining the relative importance of factors, we hope to provide a basis for developing recommendations for building or maintaining stand and landscape resilience and resistance to intense wild re and EWE.

The survey consists of three main sections: First, some brief questions about your background, then questions about the factors that determine the resilience and resistance of stands and landscapes to EWE, and finally questions about the factors that determine post- re dynamics.

IT SHOULD NOT TAKE MORE THAN 15 MINUTES TO FILL OUT THIS SURVEY

Thank you in advance for taking the time to complete this survey.

*Required

INFORMED CONSENT*

Within FIRE-RES Project, part of the European Union's Horizon 2020 research and innovation programme under grant agreement No 101037419, you are invited to participate in the following survey on fuel-related factors, metrics and thresholds driving extreme wildfire events and post-fire dynamics. From CTFC, we thank you for your participation.

The information collected in this questionnaire is anonymous and absolutely confidential. Your name will not appear in any report or result. The results will be used for research/technical purposes only.

Your participation in this research study is voluntary. You may choose not to participate. If you decide to participate in this activity, you may withdraw at any time.

Tick all that apply.

I have understood the contents and objectives of the questionnaire and I consent to participate voluntarily.

DATA PROTECTION *

CTFC as Data controller, collects this data through Google Forms to carry out a study on experts' views on the thresholds of different fuel-related factors to prevent the development of EWE and data for Central and Northern Europe. By checking the acceptance box, you give your consent. The data will not be transferred to any country or international organization outside European Union. This information will be stored on Google's servers. You can see their privacy policy at <https://policies.google.com/privacy?hl=en> and the CTFC Privacy Policy at <https://www.ctfc.cat/en/protecciodades.php>

INFORMATION ON DATA PROTECTION

Data controller: Forest Science and Technology Centre of Catalonia (CTFC) Aim: Data collection to assess the factors, metrics and thresholds driving extreme wildfire events and post-fire dynamics.

Data processor: Google Ireland Limited.

Type of data: name and surnames, email, profession, expertise, country. Rights: Access, rectify, oppose the use, limit the use and delete your data specify in CTFC privacy policy. You can also contact us at: dpd.ctfc@ctfc.cat Duration: Your data will be stored for the time necessary to carry out the purposes for which it was collected or until you revoke your consent.

Tick all that apply.

I have read and accept the CTFC Privacy Policy.

YOUR BACKGROUND

Q1) We need basic information on your expertise.

A) Can you tell us which bioregion you consider yourself an expert? Please, * select just one.

Mark only one oval.

- Macaronesia
- Mediterranean
- Atlantic
- Alpine
- Continental
- Boreal
- Other:

B) Can you tell us what aspects of wildfires you consider yourself an expert in? *
Select as many options as you need.

Tick all that apply.

- Fire behaviour
- Fuel management
- Fire ecology
- Post- re management
- Other:

C) What is your professional position? *

Mark only one oval.

- Academic (Researcher, Post-doctoral researcher, PhD student)
- Forest manager
- Fire responder (Wildfire analyst, fire fighter)
- Other:

D) How many years of experience do you have in your position? *

E) Can you tell us in which specific country do you carry out your activity? *

FACTORS DRIVING RESISTANT AND RESILIENT LANDSCAPES TO EWE

Definitions of EWE, forest resistance and resilience are provided below to be considered while answering to the questionnaire.

EWE: wild fire with large-scale complex interactions between fire and atmosphere generating pyroconvective behaviour, coupling processes, that results in fast, intense, uncertain, and fast-paced changing fire behaviour. It results in fire behaviour exceeding the technical limits of control (fireline intensity 10.000 kW/m; rate of spread >50 m/min; spotting distance >1 km and exhibiting prolific to massive spotting based on Tedim et al. 2018, and extreme growth of rate (surface per hour, ha/h) values). At the same time, given current operational models, this extreme fire behaviour is unpredictable, with moments of observed fire behaviour well surpassing the expected. This overwhelms the decision-making capabilities from the emergency system. It may represent a heightened threat to crews, population, assets, and natural values, as well as have relevant negative socioeconomic and environmental impacts.

FOREST RESISTANCE: the ability of the ecological system to persist through the disturbance event. That is, the capacity to continue providing functions and ecosystem services immediately after the event. At the stand level, resistance could be inferred from the influence of forest structure and composition on severity and intensity. At the landscape level, resistance could be inferred from the spatial configuration and composition of patches on the rate of fire spread (Deroose and Long, 2014).

FOREST RESILIENCE: the ability of the ecological system to recover the functions and ecosystem services that the system provided before the fire. In the case of wild fire, resilience could be defined as the effect of fire on subsequent forest structure and composition (at the stand level) and on subsequent proportions of age classes and on species dominance in the landscape (at the landscape level) (Deroose & Long, 2014). Resilience depends on the characteristics of the system (e.g., diversity of plant responses to fire), the event (e.g., intensity), and the presence of additional stresses before and after the fire event (e.g., prolonged drought, pest outbreaks, torrential rains, etc.).

Tedim, F., Leone, V., Amraoui, M., Bouillon, C., Coughlan, M. R., Delogu, G. M., ... & Xanthopoulos, G. (2018). Defining extreme wild fire events: Difficulties, challenges, and impacts. Fire, 1(1), 9.

Deroose, R. J., & Long, J. N. (2014). Resistance and resilience: a conceptual framework for silviculture. Forest Science, 60 (6), 1205–1212.

POST-FIRE DYNAMICS

Previous studies have shown that fire impacts and post-fire dynamics are influenced by a number of factors related to pre-fire vegetation, fire event, landscape structure, soil properties, and topography. However, most of these studies have been conducted in fire-prone areas or are related to a specific fire event, while there is a lack of information for boreal, continental, and alpine regions.

Q9) The dropdown menu in A and B (below) includes some vegetation types from boreal, alpine, and continental bioregions for which there is not as much information on post-fire impacts and dynamics. In order to make recommendations for post-fire management and set priorities, we need to know the importance of the different factors that influence their post-fire dynamics. Select a maximum of two vegetation types (one from A and the other, if you wish, from B) for which you have more experience or knowledge, and then go to C, D, E (First vegetation type) and F, G and H (Second vegetation type).

If you select "I have limited experience or knowledge of these types of vegetation" from the BOTH dropdown menu of A and B, you can skip C, D, E, F, G, and H and submit your answers.

A) FIRST vegetation type *

Mark only one oval.

- I have limited experience or knowledge of these types of vegetation.
- Hemiboreal Mountain pine (*Pinus mugo*) forests
- Hemiboreal and continental Scots pine (*Pinus sylvestris*) forests
- Alpine Scots pine or Black pine (*Pinus nigra*) in the Alps or *Pinus uncinata* in the pyrenees forests
- Subalpine larch (*Larix sp.*) forests
- Spruce (*Picea abies*) forests
- Fir (*Abies alba*) forests
- Tall deciduous oak (*Quercus sp.*) forest
- Mixed *Quercus sp.* and *Fraxinus* forests
- Beech (*Fagus sp.*) forests
- Swiss stone pine (*Pinus cembra*) forests

B) SECOND vegetation type *

Mark only one oval.

- I have limited experience or knowledge of these types of vegetation.
- Hemiboreal Mountain pine (*Pinus mugo*) forests
- Hemiboreal and continental Scots pine (*Pinus sylvestris*) forests
- Alpine Scots pine or Black pine (*Pinus nigra*) in the Alps or *Pinus uncinata* in the pyrenees forests
- Subalpine larch (*Larix sp.*) forests
- Spruce (*Picea abies*) forests
- Fir (*Abies alba*) forests
- Tall deciduous oak (*Quercus sp.*) forest
- Mixed *Quercus sp.* and *Fraxinus* forests
- Beech (*Fagus sp.*) forests
- C) FIRST vegetation type (if selected):**

For each of the following metrics related to fire regime, pre-fire vegetation, and postfire short-term competition, give a rating from 1 to 10 according to its importance in limiting

post-fire recovery of the selected vegetation type (1 irrelevant, 10 very important; move the horizontal scroll bar to view all ratings).

Mark only one oval per row.

	I do not know	1	2	3	4	5	6	7
Fire severity	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fire frequency	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Summer fire	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Spring fire	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fall fire	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Young forest	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Mature forest	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Canopy cover (%)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Tree vigour and health	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Hervibory	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Competition with pioneer species	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Is there any comment you would like to share (e.g., any missing factor, explain in more detail a factor)?

D) FIRST vegetation type (if selected):

For each of the following metrics related to climatic and topographic factors, give a rating from 1 to 10 according to its importance in limiting post-fire recovery of the selected vegetation type (1 irrelevant, 10 very important; move the horizontal scroll bar to view all ratings).

Mark only one oval per row.

	I do not know	1	2	3	4	5	6	7	8	9	10
Pre-fire: Long drought event	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Post fire: Long drought event	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Post-fire: Torrential rain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Slope: High	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Aspect: North	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Aspect: East	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Aspect: South	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Aspect: West	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Is there any comment you would like to share (e.g., any missing factor, explain in more detail a factor)?

E) FIRST vegetation type (if selected):

For each of the following metrics related to soil characteristics, give a rating from 1 to 10 according to its importance in limiting post-fire recovery of the selected vegetation type (1 irrelevant, 10 very important; move the horizontal scroll bar to view all ratings).

Mark only one oval per row.

	I do not know	1	2	3	4	5	6	7	
pH acidic	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
High erosion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Anoxic soil	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Stoniness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Soil depth	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Sandy soil	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Clay soil	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(

←
→

Is there any comment you would like to share (e.g., any missing factor, explain in more detail a factor)?

F) SECOND vegetation type (if selected):

For each of the following metrics related to fire regime, pre-fire vegetation, and postfire short-term competition, give a rating from 1 to 10 according to its importance in limiting post-fire recovery of the selected vegetation type (1 irrelevant, 10 very important; move the horizontal scroll bar to view all ratings).

D1.12 INNOVATIVE POST-FIRE STRATEGIES AND ADAPTATION TO THE CURRENT CONTEXT OF INCREASING ENVIRONMENTAL UNCERTAINTIES

Mark only one oval per row.

	I do not know	1	2	3	4	5	6	7
Fire severity	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fire frequency	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Summer fire	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Spring fire	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fall fire	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Young forest	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Mature forest	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Canopy cover (%)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Tree vigour and health	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Hervibory	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Competition with pioneer species	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Is there any comment you would like to share (e.g., any missing factor, explain in more detail a factor)?

G) SECOND vegetation type (if selected):

For each of the following metrics related to climatic and topographic factors, give a rating from 1 to 10 according to its importance in limiting post-fire recovery of the selected vegetation type (1 irrelevant, 10 very important; move the horizontal scroll bar to view all ratings).

Mark only one oval per row.

	I do not know	1	2	3	4	5	6	7	(
Pre-fire: Long drought event	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Post fire: Long drought event	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Post-fire: Torrential rain	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Slope: High	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Aspect: North	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Aspect: East	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Aspect: South	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(
Aspect: West	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	(

Is there any comment you would like to share (e.g., any missing factor, explain in more detail a factor)?

H) SECOND vegetation type (if selected):


For each of the following metrics related to soil characteristics, give a rating from 1 to 10 according to its importance in limiting post-fire recovery of the selected vegetation type (1 irrelevant, 10 very important; move the horizontal scroll bar to view all ratings).

D1.12 INNOVATIVE POST-FIRE STRATEGIES AND ADAPTATION TO THE CURRENT CONTEXT OF INCREASING ENVIRONMENTAL UNCERTAINTIES

Mark only one oval per row.

	I do not know	1	2	3	4	5	6	7
pH acidic	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
High erosion	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Anoxic soil	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Stoniness	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Soil depth	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Sandy soil	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Clay soil	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Is there any comment you would like to share (e.g., any missing factor, explain in more detail a factor)?



FIRE-RES